



LIQUID Systemic Fungicide

- Works up to 4 weeks on certain diseases.
- Not affected by rain after spray has dried.
- Prevents and stops major diseases on roses, flowers, lawns, trees and shrubs.

KEEP OUT OF REACH OF CHILDREN
WARNING

SEE BACK PANEL AND ATTACHED BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, FIRST AID AND DIRECTIONS FOR USE.

ACTIVE INGREDIENTS:
Propiconazole: (CAS No. 60207-90-1)1.55%

OTHER INGREDIENTS:.....98.45%

TOTAL:100.00%

Net Contents 8 FL. OZS.

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

If ingested, induce emesis or lavage stomach. Treat symptomatically. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call **1-800-888-8372**.



230 FM 87 • BONHAM, TX 75418
EPA Reg. No. 100-773-7401
EPA Est. No. 7401-TX-1
U.S. Patent No.: 4,079,062
Visit Us At: www.fertilome.com



PEEL FROM CORNER OF BOOKLET

20M-11-210M

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Causes substantial, but temporary eye injury. Do not get in eyes. Wear goggles or safety glasses. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow people or pets to contact treated areas until sprays are dry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STOP! READ ALL INSTRUCTIONS INSIDE BEFORE USE. FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS.

GENERAL INFORMATION

This water-based Systemic Fungicide effectively prevents and stops major Diseases on Roses, Flowers, Lawns, Trees and Shrubs.

USE TIPS

- This product is highly effective when it is used to prevent Diseases or applied at the first sign of Disease. Consider treating early in the Season. This product will not eliminate Disease already present.
- When used as directed, this product is safe to Plants and controls the Diseases listed. Plants not listed may be injured by this product and Diseases not listed may not be controlled.
- Do not apply this product to African Violets, Begonias, Boston Ferns, or Geraniums.
- If you need more information about Lawn and Ornamental Diseases, contact your local County Extension Service.

MIXING PROCEDURES

Shake well before using. Apply only with a pump-up pressure sprayer, concentrate plus water-style, or a hose-end sprayer.

Note: 1/2 Fl. oz. = 1 Tablespoon; 1 fl. oz. = 2 Tablespoons

Food utensils, such as measuring cups and spoons, must not be used for food purposes after use in measuring pesticides.

PUMP-UP SPRAYER:

1. Fill with required amount of water.
2. Add the amount of concentrate specified in the charts that follow.
3. Shake well before application.

CONCENTRATE PLUS WATER-STYLE SPRAYER:

1. From the following chart, determine the amount of concentrate required.
2. Pour concentrate into sprayer jar to the required "fl. oz." level.
3. Then add the required amount of water to the "gals." level.
4. Replace sprayer top on jar and shake well before spraying.

DIAL-STYLE HOSE-END SPRAYER:

1. Set sprayer on the "fl. oz./gallon" level specified in the following chart.
2. Use only enough product for the area to be treated.
3. Close sprayer, attach to hose and apply product to Lawn. (For example: to treat a 1,000 sq. ft. Lawn for Powdery Mildew, pour 12 fl. oz. into sprayer jar. Set sprayer dial at 3 fl. oz. per gallon.)

APPLICATION INSTRUCTIONS

STOP! DO NOT ALLOW PEOPLE OR PETS TO CONTACT TREATED AREAS UNTIL SPRAY HAS DRIED.

ROSES AND FLOWERS

- General Directions:**
- 1) Apply product to the point just when it begins to run off of the leaves.
 - 2) Allow spray to completely dry before watering.

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Roses	Powdery Mildew	1 Tablespoon (1/2 fl. oz.) per gal. of water	Thorough coverage needed for best results. Begin application at bud break. Treat every 10 days.
	Black Spot, Rust	2 Tablespoons (1 fl. oz.) per gal. of water	Treat every 10 days. Note: May not provide 100% control of Black Spot under severe conditions.
Calendula	Powdery Mildew	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 30 days.
Carnation, Iris	Rust	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 10 to 14 days.
Chrysanthemums	Ray Blight	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 7 days.
Delphinium	Powdery Mildew	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 14 days.

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Marigold	Leaf Blight, Leaf Spots	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 14 days.
Phlox, Zinnia	Powdery Mildew	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 10 to 14 days.
Snap- dragons	Rust	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 10 days.
Sweet William	Leafspot	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 14 days.

LAWNS

- 1) Apply after mowing or allow sprayed area to completely dry before mowing.
- 2) Measure the length and width of area to be treated before application.
Length x width = square feet. For example, an area 25 ft. long and 20 ft. wide is 500 sq. ft.

ESTABLISHED LAWN

DISEASE	AMOUNT OF PRODUCT	COMMENTS
Brown Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	If needed, repeat treatment every 10 days. First application should be in May or June before disease is present.
Dollar Spot, Powdery Mildew, Rust, Anthracnose	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply every 14 days.
Fusarium Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	If needed, repeat every 21 days when conditions are favorable for disease to occur.
Necrotic Ring Spot	9 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply in the fall and/or the early spring.
Red Thread, Pink Patch, Dichondra Rust on Dichondra Lawns	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	If needed, repeat every 21 days when conditions are favorable for disease to occur.
Snowmold (Gray or Pink)	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply twice in the fall prior to snow cover waiting 14 days between each application.
Spring Dead Spot	9 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply when conditions are favorable for disease development.

ESTABLISHED LAWNs (Continued)

DISEASE	AMOUNT OF PRODUCT	COMMENTS
Stripe Smut	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply in Fall after grass is dormant or in early Spring before grass starts to grow.
Take-All Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply twice in the fall (Sept. and Oct.) and twice in the spring (April and May). Southern lawns also need to be treated twice in the summer (June - August).
Zoysia Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Make 1 application in the early fall prior to the development of disease symptoms.

NEW SEEDLINGS

AMOUNT OF PRODUCT	COMMENTS
5 fl. oz. in 2 gals. of water per 500 sq. ft. of lawn	Apply at the 2 to 3-leaf stage of growth for faster root development and top growth.

TREES AND SHRUBS

- General Directions:** 1) Spray just until product runs off leaves.
2) Allow the spray to dry before watering plants.

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Ash	Powdery Mildew	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	Apply every 21 days.
Azalea	Flower blight	1/2 Tablespoon (1/4 fl. oz.) per gal. of water	Apply when there is a 5% bloom and again at 70% bloom.
Crab Apple Do not use fruit from treated crab apples for food purposes.	Scab, Rust	1/2 Tablespoon (1/4 fl. oz.)	Apply when buds just begin to break open. For scab, apply every 14 days. For rust, make 3 applications, beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 1/2 Tablespoons (3/4 fl. oz.) per gal. of water	Apply every 14 days.
Crape Myrtle	Powdery Mildew	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	Apply every 21 days.

TREES AND SHRUBS *(Continued)*

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Dogwood	Anthrax-nose	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 14–28 days.
Douglas Fir, Shasta Fir	Needle Rust	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	For Douglas Fir, apply every 21 days. For Shasta Fir, apply in May–June before disease occurs
Hawthorn, Poplars	Rust	1/2 Tablespoon (1/4 fl. oz.) per gal. of water	Apply every 14-21 days.
Jeffrey Pine	Tip Blight	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	Make the first application in mid–late winter. (Nov.–Jan.) Apply every 2 months up to 3 times.
Juniper	Blight	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 14-21 days.
Lilac	Powdery Mildew	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	Apply every 21 days.

TREES AND SHRUBS (Continued)

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Oak Red Tip Photinia	Leaf Spot	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	Apply every 14 days.
Pines	Conifer Blights	3 Tablespoons (1 1/2 fl. oz.) per gal. of water	Apply every 14 to 21 days.
Pyra- can- tha	Leaf Spot	1/2 Tablespoon (1/4 fl. oz.) per gal. of water	Apply every 14 to 21 days.
Rhodo- dendron	Flower Blight	1/2 Tablespoon (1/4 fl. oz.) per gal. of water	Apply when there is a 5% bloom and again at 70% bloom.

NON-BEARING FRUIT AND NUT TREES

- General Directions:** 1) Non-bearing fruit and nut trees are those trees that will not produce fruit for at least one year after application of this product.
2) Apply product to the point just when it begins to run off of the leaves.

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Apple	Scab	1/2 Tablespoon (1/4 fl. oz.) per gal. of water	Start applications when leaves start growing in the Spring and spray every 14 days.
	Rust	1/2 Tablespoon (1/4 fl. oz.) per gal. of water	Make 3 applications beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 Tablespoon (1/2 fl. oz.) per gal. of water	Start applications in early Spring and spray every 14 days.
Citrus	Greasy Spot	4 Tablespoons (2 fl. oz.) per gal. of water	Apply June–August every 21 days.

NON-BEARING FRUIT AND NUT TREES (Continued)

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Pecan	Scab, Powdery Mildew, Downy Spot, Zonate Leaf, Fungal Leaf Scorch, Brown Spot	2 Tablespoons (1 fl. oz.) per gal. of water	Start applications at bud break and make 3 applications at 14-day intervals.
Walnut	Anthraco- nose	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 14 days.

BEARING FRUIT AND NUT TREES

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Apricots, Cherries, Nectarines, Peaches, Plums and Prunes Do not apply to "Stanley" type plums	Brown Rot, Blossom blight, Powdery Mildew, Cherry Leaf Spot, and Fruit Brown Rot	2 Tablespoons (1 fl. oz.) per gal. water	Start applications in early Spring and spray every 21 days, making no more than 4 applications

SWEET CORN

PLANT	DISEASE	AMOUNT OF PRODUCT	COMMENTS
Sweet Corn	Leaf blights, Rust, Gray Leaf Spot, and Eye Spot	2 Tablespoons (1 fl. oz.) per gal. water	Apply when disease first appears and continue every 14 days, making no more than 4 applications.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place, away from heat or open flame.

PESTICIDE DISPOSAL: **If empty:** Do not reuse container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness of other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of VOLUNTARY PURCHASING GROUPS, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold VOLUNTARY PURCHASING GROUPS, Inc. and Seller harmless for any claims relating to such factors.

VOLUNTARY PURCHASING GROUPS, Inc. warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or VOLUNTARY

PURCHASING GROUPS, Inc., and Buyer and User assume the risk of any such use. VOLUNTARY PURCHASING GROUPS, Inc. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall VOLUNTARY PURCHASING GROUPS, Inc. or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF VOLUNTARY PURCHASING GROUPS, Inc. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES, OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF VOLUNTARY PURCHASING GROUPS, Inc. OR SELLER, THE REPLACEMENT OF TH PRODUCT.

VOLUNTARY PURCHASING GROUPS, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions for Sale and Limitation of Warranty, which may not be modified except by written agreement signed by a duly authorized representative of VOLUNTARY PURCHASING GROUPS, Inc.

Local Solutions For Local Problems™



For More Lawn and Garden Products Visit fertilome.com



Liquid Systemic Fungicide

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 100-773-7401

EPA Est. No. 7401-TX-1

U.S. Patent No.: 4,079,062

Visit Us At: www.fertilome.com

