



BEFORE

AFTER

ferti·lome®

**WEED ZONE  
FREE**

Ready To Use

- Rain-fast in as little as 3 hours • Visible results in just 24 hours
- Proven cool-weather performance

TRIMEC® is a registered trademark of PBI/Gordon Corporation

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

SEE BACK PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

**ACTIVE INGREDIENTS:**

MCPA, 2-ethylhexyl ester .....	0.3370%
Mecoprop-p acid .....	0.0660%
Dicamba acid .....	0.0180%
Carfentrazone-ethyl .....	0.0023%

**INERT INGREDIENTS:** ..... 99.5767%

**TOTAL:** ..... 100.0000%

THIS PRODUCT CONTAINS: 2-methyl-4-chlorophenoxyacetic acid  
equivalent, 0.216%

**NET CONTENTS ONE QUART (32 FL. OZ. / 946ML)**

ferti•lome® Weed Free Zone Ready-To-Use

#### FIRST AID

<b>If In Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first five minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If On Skin or Clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact  ChemTel  
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### ENVIRONMENTAL HAZARDS

This pesticide may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 48 hours.

PEEL FROM CORNER OF BOOKLET



Manufactured By:  
*Voluntary  
Purchasing  
Groups, Inc.*



230 FM 87 • BONHAM, TEXAS 75418  
EPA Reg. No. 2217-865-7401  
EPA Est. No. 7401-TX-01  
Visit Us At: [www.fertilome.com](http://www.fertilome.com)  
Product Questions?  
855-270-4776



**STOP!** READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

##### Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.

##### PRODUCT FACTS:

**ferti•lome® WEED FREE ZONE Ready-To-Use** contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

**ferti•lome® WEED FREE ZONE Ready-To-Use** offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in Turfgrass, e.g., Spurge, Pennywort (Dollarweed), Dandelion, and White Clover.
- Injury to weeds can be noticed within hours of application and plant death can occur within 7 - 14 days.

#### WHERE TO USE

**On Lawns:** Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as:

Kentucky Bluegrass	Common Bermudagrass	Hybrid Bermudagrass
Perennial Ryegrass	Mixtures of cool season species	Red or Fine Leaf Fescues
Tall Fescue	St. Augustinegrass (excluding 'Floratum')	Zoysiagrass

##### Prohibitions:

- Do not apply this product to Bentgrass, Carpetgrass, Dichondra, legumes and lawns where desirable clovers are present.
- Do not apply to 'Floratum' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.
- Do not apply more than 1.9 gallons of product per 1,000 sq. ft. per year (1.5 lb. a. i. MCPA per acre per year).
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

#### APPLICATION SCHEDULES (WHEN TO APPLY)

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant Bermudagrass, Zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

#### HOW TO USE

**ferti•lome® WEED FREE ZONE Ready-To-Use** is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide the uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

#### BROADLEAF WEEDS CONTROLLED

**ferti•lome® WEED FREE ZONE Ready-To-Use** will control or suppress the following broadleaf weeds:

#### BROADLEAF WEEDS

Aster, White Heath & White Prairie Bedstraw Beggarweed, Creeping Bindweed Black Medic Broadleaf Plantain Buckhorn Plantain Bull Thistle Burclover Burdock, Common Buttercup, Creeping Carpetweed Chickweed, Common Chicory Cinquefoil Clover Compassplant Curly Dock Dandelion Dayflower Deadnettle Dock Dogfennel English Daisy	False Dandelion (*Spotted Catsear & Common Catsear) Field Bindweed (*Morningglory & Creeping Jenny) Field Oxeye-Daisy (*Creeping Oxeye) Filaree, Whistestem & Redstem Florida Pusley Ground Ivy Groundsel Hawkweed Healall Henbit Innocence (Blue-Eyed Mary) Lambsquarters Lawn Burweed Lespedeza, Common Mallow, Common Matchweed Mouseear Chickweed Old World Diamond Flower Oxalis (*Yellow Woodsorrel & Creeping Woodsorrel)
---	--

\*Synonyms

**BROADLEAF WEEDS (Continued)**

Parsley-Piert	Spotted Spurge
Pennsylvania Smartweed	Spurge, Prostrate
Pennywort (*Dollarweed)	Thistle
Pepperweed	Veronica (*Corn Speedwell)
Pigweed	Virginia Buttonweed
Pineappleweed	White Clover (*Dutch Clover
Plantain	Honeysuckle Clover, White
Poison Ivy	Trefoil, & Purplewort)
Poison Oak	Wild Carrot
Prostrate Knotweed	Wild Garlic
(* Knotweed)	Wild Geranium
Puncturevine	Wild Lettuce
Purple Cudweed	Wild Mustard
Purslane	Wild Onion
Ragweed	Wild Strawberry
Redweed	Wild Violet
Red Sorrel (*Sheep Sorrel)	Yarrow
Shepherdspurse	Yellow Rocket

\*Synonyms

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in an area inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

The Manufacturer warrants only that the chemical composition of this product conforms to the Ingredient Statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions For Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss or profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



BONHAM, TEXAS 75418  
EPA Reg. No. 2217-865-7401  
EPA Est. No. 7401-TX-01  
Visit Us At: [www.fertilome.com](http://www.fertilome.com)  
Product Questions? 855-270-4776

10528-1114-CL

