



ferti·lome®



Crabgrass® Preventer Plus Lawn Food

Contains Team*- Granular Fertilizer

**In Established Turfgrasses preemergence control -
crabgrass, foxtail, goosegrass, poa annua &
other annual weed grasses.**

ACTIVE INGREDIENTS:
 benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine..... 0.76%
 trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 0.39%
 INERT INGREDIENTS:..... 98.85%
 TOTAL:..... 100.00%
 Contains 0.23 Pounds Active Ingredient Per 20 Pound Bag

*Trademark of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR FIRST AID, PRECAUTIONARY STATEMENTS, AND DIRECTIONS FOR USE.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies at end of label. If terms are unacceptable, return at once unopened. Do Not Ship Or Store With Food, Feeds, Drugs Or Clothing.

Net Weight 20 LBS. (9.08 KG)

Crabgrass Preventer Plus Lawn Food

Contains Team*- Granular Fertilizer

**In Established Turfgrasses preemergence control -
crabgrass, foxtail, goosegrass, poa annua &
other annual weed grasses.**

ACTIVE INGREDIENTS:
benefin: N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine..... 0.76%
trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 0.39%
INERT INGREDIENTS:..... 98.85%
TOTAL:..... 100.00%

Contains 0.23 Pounds Active Ingredient Per 20 Pound Bag

*Trademark of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR FIRST AID, PRECAUTIONARY STATEMENTS, AND DIRECTIONS FOR USE.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies at end of label. If terms are unacceptable, return at once unopened. Do Not Ship Or Store With Food, Feeds, Drugs Or Clothing.

Net Weight 20 LBS. (9.08 KG)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN**

WARNING

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. HARMFUL IF ABSORBED THROUGH SKIN. PROLONGED OR FREQUENTLY REPEATED SKIN CONTACT MAY CAUSE ALLERGIC REACTIONS IN SOME INDIVIDUALS.

Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)
ALL LOADERS, APPLICATORS, AND OTHER HANDLERS MUST WEAR:**

- Long-sleeved shirt and long pants.
- Shoes plus socks.
- Chemical-resistant gloves made of any waterproof material.
- Goggles or faceshield.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If In Eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

If On Skin Or Clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency contact **ChemTel INC.**
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585

ENVIRONMENTAL HAZARDS

Reseeding: Do not reseed treated areas for minimum of 8 weeks after single application of 174 lbs. of product per acre, or 12 to 16 weeks after a single application of 260 lbs. of product per acre unless injury can be tolerated. For split applications, do not reseed treated areas for 8 to 10 weeks after the last 130 lbs. of product per acre rate is applied, or 10 to 12 weeks after the last 174 lbs. of product per acre rate is applied unless injury can be tolerated.

**THIS PRODUCT IS RECOMMENDED FOR USE ON
THE FOLLOWING ESTABLISHED TURFGRASS SPECIES:**

Established Cool Season Turfgrasses	Established Warm Season Turfgrasses
<ul style="list-style-type: none"> • Bentgrass Fairways • Fescues • Kentucky Bluegrass • Perennial Ryegrass 	<ul style="list-style-type: none"> • Bahiagrass • Bermudagrass • Centipedegrass • St. Augustinegrass • Tall Fescue • Zoysiagrass

APPLICATION DIRECTIONS

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid slips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

**RATES, FREQUENCY and TIMING OF APPLICATION
BROADCAST APPLICATION RATES**

	Cool Season Turfgrasses	Warm Season Turfgrasses
Summer Annual Grass Weeds (Spring Application)	130 to 200 lbs./acre (3 to 4 lbs./1,000 sq. ft.)	260 lbs./acre (6 lbs./1,000 sq. ft.)
Winter Annual Grass Weeds (Fall Application)	130 to 174 lbs./acre (3 to 4 lbs./1,000 sq. ft.)	260 lbs./acre (6 lbs./1,000 sq. ft.)

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application. Apply this product 1 to 2 weeks prior to the germination of annual grass weeds since it does not control established weeds. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor.

In areas where germination of the annual weed grass species continues for extended periods of time, make a second application at the recommended rate of 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW
TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING THIS PRODUCT.**

Application Restrictions: Do not apply this product in a way that will contact any person or pet, wither directly or through drift. Keep pets out of the area during application. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter treated areas until dusts have settled. If soil incorporation is required after application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If incorporation is accomplished by watering-in, except for those people involved in the watering-in, do not enter or allow others to enter the treated area until the watering-in is complete the surface is dry after watering-in.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

GENERAL INFORMATION

This product is a granular fertilizer containing Team*, a preemergence herbicide for application to established turfgrasses for the control of the Annual Bluegrass (*Poa annua*); Smooth and Hairy Crabgrass, Goosegrass (Silver Crabgrass or Crowfootgrass), Barnyardgrass (Watergrass) and Green and Yellow Foxtail. This product also provides partial control of Prostrate Spurge (*Euphorbia humistrata*), Spotted Spurge (*Euphorbia Maculata*), Creeping Spurge (*Euphorbia serpens*), Creeping Woodsorrel (*Oxalis corniculata*) and Yellow Woodsorrel (*Oxalis stricta*). This product will not harm most nearby ornamental trees and shrubs.

second application at the recommended rate of 8 to 10 weeks after the first application on cool season turfgrass and 10 to 12 weeks after the first application on warm season turfgrass.

This product should be applied in split applications. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. For a split application, apply this product at 130 lbs. plus 130 lbs. per acre for cool season turfgrass and in the northern half of the warm season turfgrass area or 174 lbs. plus 174 lbs. per acre in the southern half of the warm season turfgrass area. Make the second application 8 to 10 weeks following the 130 lbs. per acre rate and 10 to 12 weeks following the 174 lbs. per acre rate.

SPREADER	SPREADER SETTINGS		
	Lbs. of product per 1,000 sq. ft.		
	3 lbs.	4 lbs.	6 lbs.
Scotts/Republic Accugreen (Drop)	3 1/4	4 1/4	7
Scotts/Republic Speedy Green (Broadcast)	3	3 3/4	6
ferti-lome® /Earthway Ev-N-Spred (Broadcast)	13	14	16

20-0-3

GUARANTEED ANALYSIS

Total Nitrogen (N).....	20%
0.50% Ammoniacal Nitrogen	
19.50% Urea Nitrogen	
Soluble Potash (K ₂ O).....	3%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
0.05 Water Soluble Copper (Cu)	
Iron (Fe).....	0.10%
0.05 Water Soluble Iron (Fe)	
Manganese (Mn).....	0.05%
0.05 Water Soluble Manganese (Mn)	
Zinc (Zn).....	0.05%
0.05 Water Soluble Zinc (Zn)	

Derived From: Sulfate Ammonia, Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, and Zinc Sulfate.

F370

WARRANTY DISCLAIMER

Voluntary Purchasing Groups, Inc. warrants that this product conforms to the chemical description on the label and

partial control on 1-flowered Spurge (*Euphorbia humifusa*), Spotted Spurge (*Euphorbia maculata*), Creeping Spurge (*Euphorbia serpens*), Creeping Woodsorrel (*Oxalis corniculata*) and Yellow Woodsorrel (*Oxalis stricta*). This product will not harm most nearby ornamental trees and shrubs.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not apply this product to golf course putting greens.
- Do not apply to dichondra.
- Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until turf is well established.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not make more than 2 applications of this product per year.
- **Maximum Single Application Rate:** Cool season turf — 200 lb of Turf Fertilizer — Contains Team 1.15% per acre; warm season turf — 260 lb of Turf Fertilizer — Contains Team 1.15% per acre.
- **Maximum Yearly Rate:** Cool season turf — 400 lb of Turf Fertilizer — Contains Team 1.15% per acre; warm season turf — 520 lb of Turf Fertilizer — Contains Team 1.15% per acre.
- **Minimum Re-Treatment Interval:** Cool season turf — 56 to 70 days; warm season turf — 70 to 84 days.
- This product must be watered in as soon as possible after application.

Manufactured By:



230 FM 87
BONHAM, TEXAS 75418

Visit Us At: www.fertilome.com

EPA Reg. No. 62719-150-7401 • EPA Est. No. 7401-TX-1

10907

*Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

WARRANTY DISCLAIMER

Voluntary Purchasing Groups, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in strict accordance with the Directions, subject to the inherent risks set forth below. VOLUNTARY PURCHASING GROUPS, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.


INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with the use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.) abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner or application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy of losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to at Voluntary Purchasing Groups, Inc. election, one of the following: (1) Refund of purchase price paid by Buyer or User for product brought, or (2) Replacement or amount of product used. Voluntary Purchasing Groups, Inc. shall not be liable for losses or damages, resulting from the handling or use of this product unless Voluntary Purchasing Groups, Inc. is promptly notified of such loss or damage in writing, in no case shall Voluntary Purchasing Groups, Inc. be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.



15M-2-