

ACTIVE INGREDIENT:0.160% OTHER INGREDIENTS: 98.494% THIS PRODUCT CONTAINS: 14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4) 3.34 lb (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167% (CAS 16484-77-8) 1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9) 3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8) Isomer Specific By AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

CAUTION

TO, COVERS UN TO SQUARE REEP **SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY** STATEMENTS, FIRST AID, AND DIRECTIONS FOR USE.

New York State: Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year.

NET WEIGHT 26 LBS. (11.8 KG)



Pro Turf

@



KEEP OUT OF REACH OF CHILDREN

weed Out® Pro Turf



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID		
If In Eyes:	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.	
If Swallowed:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.	
If On Skin Or Clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.	

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- · Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.

PRODUCT DESCRIPTION

This product is formulated to provide control of most annual grasses and certain broadleaf weeds

This product contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent broadleaf weed killer.

- The preemergent herbicide can be applied to established turfgrass to control crabgrass, goosegrass, foxtails, and other undesirable annual grasses.
- The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake • The post emergent broadleaf herbicide will control over 250 lawn weeds including dandelion, chickweed,
 - thistle, and clover. A partial list of broadleaf weeds is presented on this label.

WHERE TO USE

This product may be used on the following established turf species:

ESTABLISHED COOL SEASON TURFGRASS	ESTABLISHED WARM SEASON TURFGRASS	
Kentucky Bluegrass	Bahiagrass	
Perennial Ryegrass	Bermudagrass	
Fine Fescues	Buffalograss	
Tall Fescue	Zoysiagrass	

Do not use this product on dichondra, carpetgrass, or colonial bentgrass. Do not use this product on St. Augustinegrass in Florida. Do not use this product on turfgrass where desirable clovers are present. **GUIDELINES TO FOLLOW:**

The use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

 • Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- · Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays,
- Do not apply when windy.
- Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. • Do not exceed specified dosage for any area.
- Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees.

When applied as directed, this product will control and/or suppress* the following weeds:	WEEDS CONTROLLED					
Annual bluegrass* (Poa annua) Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Barnyardgrass* Annual ryegrass Bustard, indian Mustard, indian Mustar	When applied as directed, this product will control and/or suppress* the following weeds:					
Alisting of established broadleaf weeds: Amaranth, purple Amaranth, slender Aster, white heath Mustard, tumble Aster, tumble Aster, white heath Mustard, tumble Aster, tumble Aster, white heath Mustard, tumble Aster, tumble Mustard, tumble Aster, the heath Mustard, tumble Aster, tumble Aster, white heath Mustard, timble Mustard, tumble Aster, white heath Mustard, timble Mustard, timbl	Annual Grass Weeds					
Amaranth, purple Amaranth, purple Amaranth, purple Amaranth, slender Aster, heath Aster, white prairie Deadhedtle Aster, white heath Aster, white Mustard, tmble Mustard, tmble Mustard, tmble Mustard, tmble Mustard, umsay Mustard, wormsed Nettle, wood Austrid, wormsed Nettle, stinging (burning) Nettle, tall Nettle, tall Nettle, tall Nettle, tall Nettle, tall Nettle, tall Nettle, wood Penprocress (frenchweed) Penprocress (frenchweed) Penprocress (frenchweed) Pepperweed, perennial Pepperweed, pre	ı		Foxtail (green and			
Aster, heath Aster, white prairie Artichoke Austrian fieldcress Bedstraw, catchweed Bedstraw, smooth Bedstraw, pineywoods Beggarticks, hairy Betony, Florida Bindweed, Field Bindweed, Field Bittervess, hairy Bittervess, hairy Bittervess, hairy Bittervess, hairy Bittervess, hairy Bittervess, hairy Bitterved Black-eyed Susan Black medic Blood flower milkweed Buckhorn Bristly oxtongue Broomweed Buckhorn Bullnettle Burclover Burclover Burdnock Burneved, Jawn (spurweed) Burterup, creeping Buttonweed, common (poorjoe) Buttonweed, common (poorjoe) Buttonweed, common Chickweed, mouseear Hawkweed, weed Burterup, creeping Buttonweed, common Chickweed, common Chickweed, mouseear Hawkweed, weed Cathip Catisear, spotted (false dandelion) Chamber bitter Carps, make and weed Cathip Catisear, spotted (false dandelion) Chamber bitter Chickweed, common Chickweed, mouseear Chickweed, sticky Chicory Cinquefoil Clover, rabbit foot Mallow, dwarf Mustard, tumble Mustard, white Mustard, tumble Mustard, white Mustard, tumble Mustard, white Mustard, tumble Mustard, white Mustard, white Mustard, tumble Mustard, white Mustard, tumble Mustard, umble Mustard, white Mustard, tumble Mustard, white Mustard, umbie Aster, white eath Nettle, stinging (purning) Nettle, stinging (purning) Nettle, stinging (burning) Nettle, stong Parsity-piert Parshey-piert Parshey-piert Parshey-piert Parshey-piert Parshey-piert Parshey-piert Parshey-piert Parshey-p	A listing of established br	A listing of established broadleaf weeds:				
Clover, strawberry Marcela Snepardspurse Wild rape Snepardspurse Wild rape Smartweed, ladysthumb Wild strawberry	Amaranth, purple Amaranth, slender Aster, heath Aster, white heath Aster, white heath Aster, white heath Aster, purple Aster, white prairie Artichoke Austrian fieldcress Bedstraw, catchweed Bedstraw, smooth Bedstraw, Northern Bedstraw, pineywoods Beggarticks, hairy Betony, Florida Bindweed, Field Bindweed, Field Bindweed, Field Bindweed, Field Bindweed, Field Bindweed, Bedge Bitter wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Blood flower milkweed Blue lettuce Brassbuttons Bristly oxtongue Broomweed Buckhorn Bullnettle Burclover Burdock Burrweed, lawn (spurweed) Buttercup, creeping Buttonweed, virginia Carolina geranium (wild) Carpetweed Catnip Catsear, spotted (false dandelion) Chamber bitter Chickweed, common Chickweed, sticky Chicory Cinquefoil Clover, crimson Clover, hop	Daisy, oxeye Dandelion, common Dandelion, false Day flower Deadnettle Dichondra Dock, broadleaf Dock, broadleaf Dock, curly Dogbane Dogfennel Dollarweed (Pennywort) Dove weed Elderberry Falseflax False sunflower Fildelneck Fildree, redstem Filaree, redstem Filaree, whitestem Fleabane, daisy (annual) Fleabane, daisy (annual) Fleabane, singulfower Goldenrod Ground ivy Groundsel Gurmweed Hairy fleabane Hawkweed, meadow Hawkweed, meadow Hawkweed, mrange Hawkweed, orange Hawkweed, orange Hawkweed, yellow Healall Heartleaf drymary Henbit Hoary cress Horsenettle Horseweed Johnny-Jumpup violet Knawel Knotweed, prostrate Kochia Lambsquarters Lespedeza Mallow, alkali Mallow, bristly Mallow, common Mallow, wenice	Mustard, tansy Mustard, tumble Mustard, white Mustard, wormseed Nettle, stinging (burning) Nettle, tall Nettle, wood Oxalis Parsley-piert Parsnip Pearlwort Pennycress (frenchweed) Pepperweed, perennial Pepperweed, Viriginia (Peppergrass) Pigweed, prostrate Pigweed, edroot Pigweed, Russian Pigweed, smooth Pigweed, smooth Pigweed, smooth Pigweed, tumble Pineapple weed Plainis coreopsis (tickseed) Plantain, bracted Plantain, bracted Plantain, broadleaf Plantain, broadleaf Plantain, hoary Plantain, narrowleaf Plantain, slender Plantain, slender Plantain, slender Plantain, slender Plantain, slender Plantain, slender Plantain, woolly Poison oak Pokeweed Prairie sunflower Prickly lettuce (compass plant) Prickly sida Prostrate knotweed Prusley, brazil Pusley, Iarge flower Ragweed, bur Ragweed, common Ragweed, lancleaf Ragweed, western Radveed, western Radveed, western Radved, cinquefoil	Speedwell, persian Speedwell, slender Spiny amaranth Spurge, prostrate Spurweed Star of Bethlehem Strawberry, India mock Tansy ragwort Tanweed Thistle, bull Thistle, canada Thistle, bull Thistle, canada Thistle, platte Thistle, platte Thistle, platte Thistle, plumeless Thistle, plumeless Thistle, plumeles Thistle, plumeles Thistle, plumeles Thistle, rouse Thistle, rouse Thistle, rouse Thistle, wavyleaf Thistle, wayleaf Thistle, yellow (horrible) Trailing crownvetch Velvetleaf Vervain, blue Vervain, blue Vervain, tall Vetch, bird Vetch, milk Vetch, milk Vetch, narrowleaf Vetch, wild Virginia creeper Western clematis Western salsify Wild aster Wild buckweat Wild garlic Wild garlic Wild marigold Wild marigold Wild onion Wild parsnip		

Clover, white Clover, white sweet Clover, white sweet Cockle Cocklebur Cocklebur, oriental Cocklebur, spiny Common mullein Corn spurry Creeping beggarweed Creeping Jenny (creeping charilie) Cudweed Cupid shaving brush Daisy, English	Matchweed Mexicanweed Moneywort Morningglory, bigroot Morningglory, ivyleaf Morningglory, mexican Morningglory, tall Morningglory, tall Morningglory, wolly Mugwort Mustard, ball Mustard, black Mustard, blue Mustard, hedge	Smartweed, pale Smartweed, pennsylvania Smooth chaff-flower Smoothdock Sorrel, red (sheep) Sowthistle, annual Sowthistle, spiny Spanish needles Speedwell, bi-lobed Speedwell, common Speedwell, corn	Wild sweet potato Wild violet Woodsorrel, creeping Woodsorrel, violet Woodsorrel, yellow (Oxalis) Woolly croton Wormseed Yarrow, common Yarrow, western Yarrow, yellow	
* For accomplete control of Parayardarase and Googgarase, two applications made at a 6 to 10 week				

For acceptable control of Barnyardgrass and Goosegrass, two applications made at a 6 to 10 week

25-0-4

GUARANTEED ANALYSIS			
Total Nitrogen (N)			
1% Ammoniacal Nitrogen			
24% Urea Nitrogen*			
Soluble Potash(K ₂ O)			
Soluble Potash(K2O) 4% Boron (B) 0.02% Copper (Cu) 0.05%			
Copper (Cu)			
0.05% Water Soluble Copper (Cu)	1		
Iron (Fe)	1		
0.10% Water Soluble Iron (Fe)	1		
Manganese (Mn)	1		
0.05% Water Soluble Manganese (Mn)			
Zinc (Zn)			
0.05% Water Soluble Zinc (Zn)	1		
Derived From: Ammonium Sulfate, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Iron Sulfate, Manganese Sulfate, Zinc Sulfate.			
*7.5% Slowly Available Nitrogen from Polymer Coated Sulfur Coated Urea F 370			

WHEN TO APPLY

This product may be applied as a single application or as split applications in the spring, early summer, or fall. For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected germination of crabgrass. When the period of germination for crabgrass and other weeds listed on this label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial

For Crabgrass and other listed grasses: This product is a preemergent herbicide to control Crabgrass (smooth, southern, and large) and other susceptible annual grasses. This product may also provide early post emergent control of Crabgrass. However, post emergent Crabgrass control will be obtained only when this product is applied before the first tiller stage of Crabgrass.

For broadleaf weed control: Best results are obtained during the spring and fall when broadleaf weeds are present, young and actively growing under adequate soil moisture and favorable temperatures (50°to 80°F).

HOW TO APPLY

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay moving for 3 days after application.

Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and ornamental shrubbery or desirable plants

HOW MUCH TO USE

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result

For Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, and Zoysiagrass: Apply 2.5 to 3.6 pounds of product per 1.000 square feet of established turfgrass

For hybrid Bermudagrass, Bahiagrass and Buffalograss: Apply 3.0 pounds of product per 1,000 square feet of established turfgrass.

New York State: Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year.

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft.

CALIBRATION INSTRUCTIONS:

- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

SUGGESTED SPREADER SETTINGS				
SPREADER	SPREADER SETTINGS Lbs. of product per 1,000 sq. ft. 2.5 3.0 3.6.			
Scotts®/Republic Accugreen (Drop)	2-3/4 3-1/4 3-3/4			
Scotts/Republic Speedy Green (Broadcast)	3 3-1/2 4			
ferti•lome®/Earthway Ev-N-Spred (Broadcast)	14 15 16			

The above spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds, or other pesticides.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

ferti•lome® and Weed Out® are registered trademarks of Voluntary Purchasing Groups, Inc.



230 FM 87 **BONHAM, TEXAS 75418** Visit Us At: www.fertilome.com

EPA Reg. No. 2217-913-7401 • EPA Est. No. 7401-TX-1 Information regarding the contents and levels of metals in this product is available on the internet at: http://www.aapfco.org/metals.htm

Product Questions? 855-270-4776

