



ferti·lome[®]

Liquid Systemic Fungicide II

Ready To Spray

**Prevents & controls major diseases on Roses,
flowers, lawns, trees & shrubs**

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See Back Panel For Additional
Precautionary Statements and First Aid

ACTIVE INGREDIENTS:

Propiconazole: (CAS No. 60207-90-1) 1.55%

OTHER INGREDIENTS:..... 98.45%

TOTAL:..... 100.00%

NET CONTENTS ONE QUART (32 FL. OZ. / 946 ML)

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact **ChemTel** 
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a Poison Control Center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 53883-184-7401

EPA Est. No. 7401-TX-1

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776



- Not affected by rain after spray has dried
- Works up to 4 weeks for certain diseases

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 53883-184-7401

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

PRODUCT QUESTIONS? 855-270-4776

11380-0215-CL



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial, but temporary eye injury. Do not get in eyes. Wear goggles or safety glasses. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow people or pets to contact treated areas until spray is dry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product must not be used in the production of agricultural commodities grown for resale.

READ ALL INSTRUCTIONS INSIDE BEFORE USE.

FOLLOW ALL DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL. POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS MAY RESULT FROM IMPROPER USE OF THIS PRODUCT.

DO NOT ALLOW PEOPLE OR PETS INTO TREATED AREAS UNTIL SPRAY HAS DRIED.

GENERAL INFORMATION

This product is a water-based systemic fungicide which effectively prevents and controls major disease on roses, flowers, lawns, trees, and shrubs. **ferti•lome® Liquid Systemic Fungicide II Ready-To-Spray** is very effective

when used for disease prevention or applied at the first sign of disease. Best results occur when treating early in the season. If disease is already present, this product will not eliminate it.

ferti•lome® Liquid Systemic Fungicide II Ready-To-Spray, when used as directed, is safe to plants and controls the diseases listed on this label. The use on plants or diseases not listed on the label may result in plant injury and poor disease control. Do not apply this product to African violets, begonias, Boston ferns, or geraniums.

Contact your local County Extension Service if you need additional information about lawn and ornamental diseases.

**READY-TO-SPRAY INSTRUCTIONS FOR USE OUTDOORS ON
ROSES, FLOWERS, HOME LAWNS, TREES AND SHRUBS**

HOW TO USE THE READY-TO-SPRAY SYSTEM

Connect

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

Spray

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small plastic tab back and turn knob clockwise to ON position.
3. Spray evenly to the area to be treated. Refer to the sight gauge clear view strip (graduated scale) on the side of the container to determine the amount of product sprayed.

Finish

1. To stop spraying, turn knob counter clockwise to OFF position.
2. Turn off water.
3. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
4. Disconnect sprayer from hose.

ROSES AND FLOWERS

- Apply product to the point of runoff.
- Before watering, allow spray to dry fully.

APPLICATION RATES:			
Plant	Disease	How Much Product To Apply	Comments
Roses	Powdery Mildew	1 Table-spoon (1/2 fl. oz.) per gal. of water	For best results, thorough coverage is needed. Begin application at bud break. Treat every 10 days.
	Black Spot, Rust	2 Table-spoons (1 fl. oz.) per gal. of water	Treat every 10 days. Note: Under severe conditions, 100% control of black spot may not be achieved.

APPLICATION RATES (Continued)

Plant	Disease	How Much Product To Apply	Comments
Calendula	Powdery Mildew	1 Table- spoon 1/2 fl. oz.) per gal. of water	Apply every 30 days.
Chrysan- themums	Ray Blight	2 Table- spoons (1 fl. oz.) per gal. of water	Apply every 7 days.
Delphinium	Powdery Mildew	1 Table- spoon (1/2 fl. oz.) per gal. of water	Apply every 14 days.
Snap Dragons	Rust	2 Table- spoons (1 fl. oz.) per gal. of water	Apply every 10 days.
Sweet William	Leaf Spot	1 Table- spoon (1/2 fl. oz.) per gal. of water	Apply every 14 days.

LAWNS

- Apply the product to the lawn after mowing or allow the area sprayed to dry completely prior to mowing.
- Prior to application, measure the length and width of area to be treated. Length x width = square feet. (For example, an area 30 ft. long and 20 ft. wide is 600 sq. ft.)

Established Lawns

APPLICATION RATES:		
Disease	How Much Product To Apply	Comments
Brown Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Repeat treatment every 10 days if necessary. First application should take place in May or June before disease is present. For southern lawns, make applications at first sign of disease in spring or fall.
Dollar Spot, Powdery Mildew, Rust, Anthracnose	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply every 14 days.

APPLICATION RATES (Continued)

Disease	How Much Product To Apply	Comments
Red Thread, Dichondra Rust on Dichondra Lawns	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Repeat every 21 days, if needed, when conditions are favorable for disease to occur.
Stripe Smut	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply in early spring before grass starts to grow or in fall after grass is dormant.
Take-All Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply twice in the fall (September – October) and twice in the spring (April – May). For southern lawns, also make applications twice in the summer (June – August).

New Seedlings

How Much Product To Apply	Comments
5 fl. oz. in 2 gals. of water per 500 sq. ft. of lawn	For faster root development and top growth, apply at the 2 to 3-leaf stage of growth.

TREES AND SHRUBS

- Apply product to the point of run off.
- Before watering plants, allow spray to fully dry.

APPLICATION RATES:

Plant	Disease	How Much Product To Apply	Comments
Crab Apple Note: Do not use fruit from treated crab apples for food purposes.	Scab, Rust	½ Table- spoon (1/4 fl. oz.) per gal. of water	Apply when buds just be- gin to break open. Scab – apply every 14 days. Rust – Make 3 applications, beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 ½ Tablespoons (3/4 fl. oz.) per gal. of water	Apply every 14 days.

APPLICATION RATES (Continued)

Plant	Disease	How Much Product To Apply	Comments
Crape Myrtle	Powdery Mildew	3 Tablespoons (1 ½ fl. oz.) per gal. of water	Apply every 21 days.
Dogwood	Anthracnose	1 Tablespoon (1/2 fl. oz.) per gal. of water	Apply every 14-28 days
Douglas Fir, Shasta Fir	Needle Rust	3 Tablespoons (1 ½ fl. oz.) per gal. of water	Douglas Fir – apply every 21 days. Shasta Fir – apply May – June before disease occurs.

APPLICATION RATES (Continued)

Plant	Disease	How Much Product To Apply	Comments
Jeffery Pine	Tip Blight	3 Tablespoons (1 ½ fl. oz.) per gal. of water	Make the first application in mid-late winter (Nov. – Jan.) Apply every 2 months up to 3 times.
Oak, Red Tip Photinia	Leaf Spot	3 Tablespoons (1 ½ fl. oz.) per gal. of water	Apply every 14 days.

NONBEARING FRUIT AND NUT TREE

- Nonbearing fruit and nut trees are those trees that will not produce fruit for at least one year after application of this product.
- Apply product to the point of run off.

APPLICATION RATES:

Plant	Disease	How Much Product To Apply	Comments
Apple	Scab	½ Table-spoon (1/4 fl. oz.) per gal. of water	Begin applications when leaves start growing in the spring and spray every 14 days.
	Rust	½ Table-spoon (1/4 fl. oz.) per gal. of water	Make 3 applications beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 Table-spoon (1/2 fl. oz.) per gal. of water	Begin applications in early spring and spray every 14 days.

APPLICATION RATES (Continued)

Plant	Disease	How Much Product To Apply	Comments
Cherry	Powdery Mildew, Leaf Spot	1 Table-spoon (1/2 fl. oz.) per gal. of water	Begin applications in early spring and spray every 21 days.
Citrus	Greasy Spot	4 Table-spoons (2 fl. oz.) per gal. of water	Apply June – August every 21 days.
Pecan	Scab, Powdery Mildew, Downy Spot, Zonate Leaf Spot, Fungal Leaf Scorch, Brown Spot	2 Table-spoons (1 fl. oz.) per gal. of water	Begin applications at bud break and make 3 applications at 14 day intervals.
Walnut	Anthrax-nose	2 Table-spoons (1 fl. oz.) per gal. of water	Apply every 14 days.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage and Spill Procedures:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available.

WARRANTY STATEMENT

Voluntary Purchasing Groups, Inc. (VPG) warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the Directions For Use. It is impossible to eliminate all risks inherently associated with use of this product.

(continued)

WARRANTY STATEMENT (Continued)

Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of VPG. To the extent allowed by law, VPG shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except as expressly provided herein, VPG makes no warranties, guarantees, or representations of any kind, whether expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at VPG's election, the replacement of this product.

Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions.

This page left blank intentionally.

This page left blank intentionally.