




ferti·lome®



# Weed & Feed With Dimension®

## Controls Weeds Before You See Them

- For use on established cool season and warm season turfgrass
- Provides up to 4 months of crabgrass control
- For control or suppression of listed grass and broadleaf weeds

**Not for use** on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purpose.

**Not for use** on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

**For use on plants** intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or on golf courses or lawns.

**In the state of New York**, do not exceed 4.6 lbs. per 1,000 sq. ft. per year of product (equivalent to 0.25 lbs. of active ingredient per acre).

**In the state of Florida**, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn / garden.

ACTIVE INGREDIENTS:  
 dithiopyr: 3,5-pyridinedicarboic acid, 2-(difluoromethyl)-  
 4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester..... 0.125%  
 OTHER INGREDIENTS: ..... 99.875%  
 TOTAL: ..... 100.00%

This product contains 0.0625 pounds of the active ingredient dithiopyr per 50 pound bag.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS, FIRST AID, AND DIRECTIONS FOR USE.

Net Weight 32 LBS. (14.5 KG)

Covers Up To  
10,000 Sq. Ft.

# Weed & Feed With Dimension®

## Controls Weeds Before You See Them

- For use on established cool season and warm season turfgrass
  - Provides up to 4 months of crabgrass control
- For control or suppression of listed grass and broadleaf weeds

**Not for use** on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purpose.

**Not for use** on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

**For use on plants** intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or on golf courses or lawns.

**In the state of New York**, do not exceed 4.6 lbs. per 1,000 sq. ft. per year of product (equivalent to 0.25 lbs. of active ingredient per acre).

**In the state of Florida**, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn / garden.

#### ACTIVE INGREDIENTS:

dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester ..... 0.125%

OTHER INGREDIENTS: ..... 99.875%

TOTAL: ..... 100.00%

This product contains 0.0625 pounds of the active ingredient dithiopyr per 50 pound bag.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS, FIRST AID, AND DIRECTIONS FOR USE.**

**Net Weight 32 LBS. (14.5 KG)**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes eye irritation. Harmful if inhaled.

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

**FIRST AID**

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In the event of a medical or chemical emergency contact ChemTel Inc. North America 1-800-255-3924

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.**

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer recycling if available.

**If partially filled:** Call you local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**USE DIRECTIONS FOR TURF**

This turf fertilizer product containing herbicide provides up to 4 months of crabgrass control.

Ferti-lome Weed & Feed With Dimension® provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination. Florida Applications: Do not exceed 4 applications per year. Do not apply more than 27.6 lb of this product/1000 sq ft per year (equivalent to 1.5 lb of active ingredient per acre).

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

**COOL SEASON GRASSES**

Bentgrass, Creeping  
Bluegrass, Kentucky  
Fescue, Fine  
Fescue, Tall  
Ryegrass, Perennial

**WARM SEASON GRASSES**


Bahiagrass  
Bermudagrass  
Buffalograss  
Carpetgrass  
Centipedegrass  
St. Augustinegrass  
Zoysiagrass

**USE PRECAUTIONS FOR TURF**

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this

**LENGTH OF CONTROL**

Ferti-lome Weed & Feed With Dimension® will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. **Refer to the map below to determine the region of the U.S. that includes your location.** Use the table below to determine the amount of Ferti-lome Weed & Feed With Dimension® to apply in your location based on the desired length of weed control. **Use "Regions" within the United States:**



□ North  
■ South

**RECOMMENDED USE RATES**

**Amount of Ferti-lome Weed & Feed With Dimension® to Apply**  
(Pounds of product to apply per 1,000 square feet of turfgrass<sup>†</sup>)

-----Length of Crabgrass Control-----

Region of U.S.	3 - 4 months
North	3.2 lb/1000 sq ft
South	4.0 - 4.7 lb/1000 sq ft <sup>††</sup>

<sup>†</sup> 1,000 square feet equals an area 100 feet x 10 feet or square area of about 32 feet on a side.

<sup>††</sup> Use the higher rate in rate range in areas of high weed pressure or for increased length of control.

**Maximum Application Rates**

- Do not apply more than 27.6 lb of this product/1000 sq ft per year (equivalent to 1.5 lb of active ingredient per acre).
- In the state of New York, do not exceed 4.6 lb per 1000 sq. ft. per year of this product (equivalent to 0.25 lb of active ingredient per acre).

**APPLICATION EQUIPMENT AND INSTRUCTIONS**

This product can be applied with a drop or broadcast (rotary) spreader. For best results, apply this product evenly and uniformly over the turfgrass (lawn) area avoiding streaking, skips or overlaps.

Before each application, determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the label states to apply 4 lbs. per 1000 sq. ft. and your area is 2000 sq. ft, then place 8 lbs. of product in the hopper and set the spreader setting at 4-¼ (per table below) if using an Accugreen drop spreader.

A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction.

**NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.**

Spreader	Spreader Settings		
	Lbs. of product per 1,000 sq. ft.		
	3.2	4.0	4.7
Scotts/Republic Accugreen (Drop)	4	4 ¼	4 ½
Scotts/Republic Speedy Green (Broadcast)	3 ¼	3 ¾	4 ¼
Ferti-lome/Earthway Ev-N-Spred (Broadcast)	14	16	17

**TERMS AND CONDITIONS OF USE**

This product should be applied only to lawns and ornamental turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. **Exception:** In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

-Use of this product on lawns and ornamental turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrass injury.

-For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. **Do not** apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

**-Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 16 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

**Other Use Precautions**

**-Do not** use clippings from treated turf for mulching around vegetables or fruit trees.

**-Do not** allow people or pets to enter treated area until dust has settled.

**APPLICATION TIMING**

**Spring Application**

**Control of Crabgrass:** Ferti-lome Weed & Feed With Dimension® provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior to their germination.

**Control or Suppression of Other Grass and Broadleaf Weeds:** For control or suppression of other grass and broadleaf weeds, this product must be applied as a **preventative** treatment prior to their germination and emergence.

**Fall Application**

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, **do not** reseed, overseed or sprig turfgrasses within 12 weeks after application.

**TIPS FOR IMPROVED WEED CONTROL**

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

**WEEDS CONTROLLED OR SUPPRESSED**

Ferti-lome Weed & Feed With Dimension will provide control of crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior to their germination.

Grasses		Broadleaf Weeds	
barley	goosegrass	bittercress †	pigweed, redroot
barnyardgrass	foxtail, green	carpetweed †	parsley-piert †
bluegrass, annual	foxtail, yellow	chickweed †	purslane, common
brome	kikuyugrass †	geranium, Carolina †	rocket, London
crabgrass, large	oats, wild	henbit	sheepspurse
crabgrass, smooth	ryegrass (annual & perennial)	knotweed, prostrate	speedwell, corn †
crabgrass, Southern	sandbur	lespedeza, common †	spurge, garden
crowfootgrass †	smutgrass	marehail	spurge, prostrate
dallisgrass		medic, black	spurge, spotted
		mustard	woodsorrel, creeping
		oxalis, buttercup	woodsorrel, yellow
		pineappleweed	

† suppression only

**TERMS AND CONDITIONS OF USE**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

**WARRANTY DISCLAIMER**

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**INHERENT RISKS OF USE**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by the buyer.

**LIMITATION OF REMEDIES**

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at the Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for the product bought, or
2. Replacement of amount of product used

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

Terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Dimension® is a registered trademark of Dow AgroSciences

**20-0-4  
GUARANTEED ANALYSIS**

Total Nitrogen (N) .....	20%
20% Urea Nitrogen*	
Soluble Potash (K <sub>2</sub> O) .....	4%
Boron (B) .....	0.02%
Copper (Cu) .....	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe) .....	0.10%
0.10% Water Soluble Iron (Fe)	
Manganese (Mn) .....	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn) .....	0.05%
0.05% Water Soluble Zinc (Zn)	

**Derived From:** Urea, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate.

\* 5.25% Slowly Available Nitrogen Derived From Polymer Coated Sulfur Coated Urea

F370

Manufactured by:



230 FM 87  
BONHAM, TEXAS 75418

Visit Us At: [www.fertilome.com](http://www.fertilome.com)

EPA Reg. No. 62719-483-7401 • EPA Est. No. 7401-TX-1

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>



Get the free mobile app at <http://gettag.mobi>

