

Do no repeat applications more often than every 7 days. The maximum number of applications per year allowed is listed in the next [].

GRAPES: (7) [5] To kill Grape Leaf Folder, Grape Leafhopper and Grape Leaf Skel-tonizer, Grape Berry Moth, Japanese Beetle and Red Banded Leaf Roller.

STRAWBERRIES: (7) [5] To kill Meadow Spittlebug, Strawberry Leaf Roller, Straw-berry Weevil, Japanese Beetles and Leafhoppers.

BLUEBERRIES: (7) [5] To kill Blueberry Maggot, Cherry and Cranberry Fruitworms and Japanese Beetle. Apply 3 weeks before harvest and repeat 10 days later if nec-essary.

CRANBERRIES: (7) [5] To kill Cutworms, Fireworms, Fruitworms, Japanese Beetle, and Leafhoppers.

CUCUMBER, MELONS, PUMPKIN AND SQUASH: (3) [6] To kill Pickleworms Mel-onworm, Cucumber Beetle, Flea Beetle, Leafhopper and Squash Bugs. May cause some leaf injury during high humidity. Do not use on watermelons in Florida.

TOMATO, EGGPLANT AND PEPPER: (3) [7] To kill Colorado Potato Beetle, Flea Beetle, Leafhopper, European Corn Borer, Fall Armyworm, Lace Bugs, Tomato Fruit-worm, Tomato Hornworm, Tarnish Plant Bugs and Stink Bugs.

OKRA: (3) [4] To kill Stink Bugs. Repeat at 7 day intervals.

CARROTS: (7) [6] To kill Flea Beetles, Leafhoppers, Six-Spotted Leafhoppers (aster yellows vector), Armyworms, Harlequin Bug, Spittlebug and Stink Bugs.

ORNAMENTAL SHRUBS, TREES AND FLOWERS

For outdoor use only. To kill Bagworm, Blister Beetles, Boxelder Bug, Boxwood Leafminer, Flea Beetles, Japanese Beetle, Lace Bugs, Leafhoppers, Leafrollers, Pe-riodical Cicada, Plant Bugs, Psyllids, Rose Aphid, Rose Slug, Scale insects in the crawler stage, Tent Caterpillars and exposed Thrips, apply thoroughly to the infested plants at the first sign of damage. For best results on scale, apply in Spring and early Summer when scale crawlers are present. Repeat applications up to 3 times per year, but not more often than once every 7 days. Do not apply to plants above chest height.

HOME LAWNS:

To kill Chinch Bugs, Sod Webworms (Lawn Moths), Fall Armyworms, Cutworms, Earwigs, Leafhoppers, and Millipedes, sprinkle lightly on grass.

To kill Ants (except harvester, pharaoh, and carpenter ants), Chiggers, Fleas, Ticks and Spiders (except Brown recluse and Black widow), sprinkle lightly on grass. Will only kill pests present at the time of application. Pest must be directly contacted by product to be effective. For best results apply after rain or watering and do not water for at least 2 days after application. Repeat treatment as needed, but not more than every 2 weeks and not more than 4 times per year.

EARWIGS

This material is effective in killing earwigs and when properly used will reduce the number gaining entrance to homes and buildings. Lightly dust a band 3 to 4 inches wide around the outside foundation wall of the home. Be sure to treat in front of all steps and other entrance areas. Earwigs breed over wide areas around yards and fields and are driven by cold weather to seek shelter in homes. Apply also around re-fuse piles, lumber, mulch and other concealment areas. Repeat treatment as needed, but not more than every 2 weeks and not more than 4 times per year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool, dry, secured area.

PESTICIDE DISPOSAL: Contact your local solid waste agency for disposal instruc-tions for unused pesticide.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this con-tainer. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS PRODUCT NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HERewith. VOLUNTARY PURCHASING GROUPS, INC. WARRANTS THIS PRODUCT TO CONFORM TO THE CHEMICAL DE-SRIPTION ON THE LABEL AND FOR THE PURPOSES STATED ON THE LABEL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE LIMIT OF ANY LIABILITY INCURRED SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER.

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 7401-69

EPA Est. No. 7401-TX-1

Visit Us At: www.hi-yield.com

50M-5-1111



Hi-Yield®

Garden Dust FOR OUTDOOR HOMEOWNERS USE



- Ready To Use 5% Dust.
- Kills Fleas and Ticks
- Kills Japanese Beetle
- Kills Listed Insects on Listed Home Garden Fruits, Vegetables, Lawns and Ornamentals

ACTIVE INGREDIENT:
Carbaryl(1-naphthyl N-methylcarbamate)..... 5.00%
OTHER INGREDIENTS:..... 95.00%
TOTAL:..... 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back/Side Panel For Additional Precautionary Statements

NET WEIGHT 1 LB. (454 g)

31120

READ ENTIRE LABEL BEFORE EACH USE

FIRST AID N-methyl carbamate

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact ChemTel
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585

NOTE TO PHYSICIANS: Carbaryl is an N-methyl carbamate that inhibits cholin-esterase. Atropine is antidotal. DO NOT use 2-Pam, opiates or cholinesterase-inhibiting drugs.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through the skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

BEE CAUTION: MAY KILL HONEY BEES AND OTHER BEES. This product is highly toxic to honey bees and other bees. Do not apply to plants in bloom.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **USE STRICTLY IN ACCORDANCE WITH LABEL RESTRICTIONS AND DI-RECTIONS.**

Restrictions

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Non-refillable container. Do not cut package to expose contents. Discard package when empty.

Precautions

Some phytotoxicity may occur on tender foliage in the presence of rain or high hu-midity of several days' duration following application.

Hi-Yield® Garden Dust is an insecticide formulation for kill of listed insects found in home gardens. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

DIRECTIONS

For residential use on listed Ornamentals, Flowers, Shrubs, and Garden Vegetables. For outdoor use only.

ORNAMENTAL FLOWERS, ORNAMENTAL SHRUBS AND ORNAMENTAL TREES to kill: Elm Leaf, Rose and Apple Aphids, Rose Slugs, Birch Leaf, Boxwood and Oak Leaf Miners, Mealy Bugs, Leaf Rollers, Plant, Lace and Boxelder Bugs, Bagworms, Japanese Beetle, Tent Caterpillars, Gypsy Moth.

Do not use on Boston Ivy as damage may result.

WHEN & HOW TO APPLY:

- Begin applications when insect or damage is first observed and repeat as directed below.
- Wearing household latex or rubber gloves, use this ready-to-use shaker can to ap-ply product. Dust lightly to cover upper leaf surfaces with a thin, even film of dust.
- Apply when air is calm.
- Start applying at the farthest corner of the treatment area and work backward. For plants above chest high, consider using a liquid carbaryl product.

The Pre-Harvest Interval (PHI) indicates the number of days between the last ap-plication and harvest. The PHI is listed in the () immediately after each crop heading.