FOR ORGANIC GARDENING

Ready to Use

ethn

Controls Blackspot, Powdery Mildew, Rust, Spider Mites, Aphids, Whiteflies and Other Insect Pests

ACTIVE INGREDIENTS:	
Clarified Hydrophobic Extract of Neem Oil	0.90%
OTHER INGREDIENTS	<u>99.10%</u>
TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN

See Back Panel For Additional Precautionary Statements.

NET CONTENTS ONE QUART (32 FL. OZ. / 946ML)

	FIRST AID
lf Inhaled:	 Move person to fresh air. If person is not breath- ing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in Eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
lf On Skin Or Clothing:	 Take off contaminated clothing. Rinsing skin immediately with plenty of water for 15-20 min- utes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In the event of a medical or chemical emergency, contact Chemital Worldwide Intl. + 01- 813-248-US88 North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-US88	



Natural Guard Brand Neem Ready-To-Use

230 FM 87 • BONHAM, TEXAS 75418 EPA Reg. No. 70051-13-7401 EPA Est. No. 7401-TX-01 Visit Us At: www.fertilome.com Product Questions? 855-270-4776



DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL INFORMATION

- Multi-purpose fungicide/insecticide/miticide 3 in 1 product.
- · Kills egg, larvae and adult stages of insects.
- Prevents fungal attack of plant tissue.
- · Not for use in food-handling areas.
- · Shake well before using.
- Thorough plant coverage is essential to obtain optimal levels of control.
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment.
- As with other pest control products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use with care on plants with tender tissue. Check for leaf burn in small-scale trials prior to use.

FUNGICIDE USE

Application Instructions

- This product is an effective fungicide for prevention and control of various fungal diseases including Black Spot on Roses, Powdery Mildew, Downy Mildew, Anthracnose, Rust Leaf Spot, Botrytis, Needle Rust, Scab, and Flower, Twig and Tip Blight.
- This product may be used on all Roses, flowers, houseplants, ornamental trees and shrubs, fruits, nuts, and vegetables grown in and around the home and home garden.
- As a preventative, apply this product on a 7 to 14 day schedule until the potential for disease development is no longer present. Spray to run off. Complete coverage of all plant tissue is necessary for control.
- To control disease already present, apply this product on a 7-day schedule until disease pressure is eliminated.

1

- Then continue spraying on 14-day schedule to prevent the disease from recurring. Spray to run off. Complete coverage of all plant tissue is necessary for control.
- To prevent Rust Leaf Spot diseases, Anthracnose, and Scab, begin applications at the first sign of spring budding.
- To prevent Powdery Mildew, apply in mid-summer or when disease is first detected. Continue applications until disease pressure no longer exists.
- To prevent Black Spot on Roses, begin applications at first bud emergence in spring. Apply this product on a 14-day preventative schedule through the spring and fall to prevent disease emergence.

INSECTICIDE USE

Application Instructions

- This product is an effective insecticide for the control of various insect pests including Whiteflies, Mealy Bugs, Armyworms, Hornworms, Leafhoppers, Bagworms, Leafminers, Psyllids, Fruit Flies, Loopers, Budworms, Beetles (Including Japanese Beetles), Aphids, Scales, Caterpillars, Midges, Budworms, and Mites (including Spider Mites).
- This product may be used on all Roses, flowers, houseplants, ornamental trees and shrubs, fruits, nuts, and vegetables grown in and around the home and home garden.
- Apply this product at first sign of insects/mites. This product is most effective in controlling insects/mites when applied on a 7-14 day schedule. For heavy insect populations, apply on a 7-day schedule. Spray to run off. Complete coverage of all plant tissue is necessary for control.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage**: Store upright at room temperature. Avoid exposure to extreme temperatures. **Pesticide Disposal and Container Handling: If empty:** Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY AND LIMITATION OF DAMAGES

Voluntary Purchasing Groups, Inc, warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the Directions For Use. Timing and method of application, weather, watering practices, nature of soil, the Insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE. Verning Burdsating 230 FM 87 • BONHAM, TEXAS 75418 EPA Reg. No. 70051-13-7401 EPA Est. No. 7401-TX-01 Visit Us At: www.fertilome.com Product Questions? 855-270-4776

Manufactured By:

40716-1114-CL

