

07778



Hi-Yield®

IMPORTED
Fire Ant Control Granules
Containing Deltamethrin



FOR RESIDENTIAL USE ONLY

- **Aids In Controlling Imported Fire Ants In Home Lawns And Other Outdoor Home Recreational Areas**

ACTIVE INGREDIENT:	
Deltamethrin: (s)-alpha-cyano-3-phenoxybenzyl-(1R,3R)-3-(2-2 dibromovinyl)-2,2-dimethyl cyclopropanecarboxylate.....	
	0.1%
INERT INGREDIENTS:	99.9%
TOTAL:	100.0%


KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements, First Aid and Directions for Use

Fire Ant Control Granules

32220

NET WEIGHT 2 LBS. (0.90 KG)



KEEP OUT OF REACH OF CHILDREN

Hi-Yield®

IMPORTED Fire Ant Control Granules

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID

If Swallowed:

- Call Poison Control Center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the Poison Control Center or doctor.
- Do not give anything by mouth to an unconscious person.

If In Eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

If On Skin Or Clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a Poison Control Center or doctor for treatment advice.

If Inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

In the event of a medical or chemical emergency contact ChemTel ChemTel North America 1-800-255-3924 or Worldwide Intl. +01-813-248-0585

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticides to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Note: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the User or Buyer.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container. Keep in an area inaccessible to children and pets.

PESTICIDE DISPOSAL: If Empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: HI-YIELD® IMPORTED FIRE ANT CONTROL GRANULES CONTAINING DELTAMETHRIN is a broad spectrum residual insecticide for the control of certain insects in home lawns.

Treat as directed-water in granules with ½ inch of water. This product is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply on saturated soil. Do not allow runoff to occur. Do not contaminate ornamental fish ponds.

In New York State, this product may not be applied to lawns that directly border coastal marshes. For control of Turf Insects other than Fleas and Ticks in New York State, a second application may be made after two weeks if target insect activity persists.

Not for use in commercial greenhouses or nurseries.

Not for use on sod farms, grass grown for seed or on golf course turf.

Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides. Do not allow children or pets on treated areas until granules have been watered in the soil and the grass or soil is dry.

HOME LAWN RECOMMENDATIONS: Use HI-YIELD® IMPORTED FIRE ANT CONTROL GRANULES CONTAINING DELTAMETHRIN to control specified pest of lawns.

Apply HI-YIELD® IMPORTED FIRE ANT CONTROL GRANULES CONTAINING DELTAMETHRIN to give complete and uniform coverage of pest infested areas. Applications should be made using the rates shown in the following chart.

PEST	DOSAGE OF PRODUCT	APPLICATION PROCEDURE
Ants, Armyworms, Bermudagrass Mites, Billbugs (Adult), Boxelder Bugs, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Digger Wasps, Essex Skipper, Field Crickets, Fiery Skipper, Fire Ants, Flea Beetles, Fleas, Grasshoppers, Hyperodes, Japanese Beetles, Lawn Moths, Leafhoppers, Mealy Bugs (Adult), Mole Crickets (aid in control), Scorpions, Silverfish and Firebrats, Sod Webworms, Sowbugs, Spiders, Spittlebugs, Sprintalis Ticks (including Lyme disease vectors), Vegetable Weevils	2 to 3 pounds per 1,000 sq. ft.	Apply this product uniformly to home lawns, under shrubbery and other ornamental recreational grassy areas. Treat when insects first appear. Immediately after application, water the lawn thoroughly with half an inch of water. Retreat only as necessary to maintain insect control
European Crane Fly Larvae (Leather Jackets)	2 to 3 pounds per 1,000 sq. ft	Apply in early April when insects are young.
Cinch Bugs	2 to 3 pounds per 1,000 sq. ft.	Start treatment when temperatures reach 80°F or above. A second application may be necessary in 6 weeks if insect activity continues.

APPLICATION EQUIPMENT: HI-YIELD® IMPORTED FIRE ANT CONTROL GRANULES CONTAINING DELTAMETHRIN is a granular formulation. Apply with a fertilizer spreader or equivalent equipment as uniformly as possible. The following table lists application equipment and recommended settings for that equipment for the application of HI-YIELD® IMPORTED FIRE ANT CONTROL GRANULES CONTAINING DELTAMETHRIN.

SPREADER SETTINGS		
SPREADER MAKE AND MODEL	SETTING TO DELIVER	
	2 lbs. per 1,000 sq. ft	3 lbs. per 1,000 sq. ft.
Scotts®/Republic Accugreen (Drop)	2 ½	3 ½
Scotts®/Republic Speedy Green (Broadcast)	2 ¾	3 ¾
ferti•lome® Earthway Ev-N-Spred (Broadcast)	13	14

NOTE: Spreader settings given are approximate and variation can occur between identical models. User should calibrate spreader by accurately weighing a few pounds into spreader and applying to a measured area. Then adjust spreader setting if necessary to ensure correct rate of application.

FLEA AND TICK CONTROL (OUTDOORS): For control of Fleas and Ticks (including, but not limited to Bear Tick, Black-legged Tick, Brown Dog Tick, Lone Star, Western Black-legged Tick, Wood Tick), a regular treatment program should be followed. Treat once a month as directed above when pests first appear in late spring (May-June) and continue until late summer (September-October).

SOD WEBWORMS: If Lawn Moths (Sod Webworms Adults) are noticed in large numbers at dusk, or at first sign of damage to lawn, treat entire lawn evenly.

PERIMETER TREATMENT: To help prevent infestation of buildings by Ants, Boxelder Bugs, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Fire Ants, Fleas, Scorpions, Sowbugs, Spiders, Springtails (Collembola) and ticks, treat a band 5 to 10 feet wide around and adjacent to the building where pests are active and may find entrance. Treat at a rate of 1 to 1½ pounds per 500 so. ft. or 2 to 3 pounds per 1,000 sp. ft.

SPOT TREATMENT: To control Ants, including Acrobat, Argentine, Big-Headed, Cornfield, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants. If possible, locate and treat nest first. Product should be applied to thoroughly cover the nest (1 to 2 teaspoons) and sprinkled with sufficient water to dissolve granules. In addition to treating Ant nests or mounds, entry points of buildings and the soil surface along the perimeter of the building should be treated.

DIGGER WASPS: Apply 1 to 2 teaspoons of granules in and around each individual ground nest opening. Make the application at dusk when Wasps are not active.

IMPORTED FIRE ANTS: Apply two tablespoons over each 8" to 10" diameter mound and up to 2 feet around to prevent escape of the queen. Immediately after application, water the granules into the soil gently with a light sprinkling can to avoid disturbing the colony. Use at least one gallon of water per mound. Apply in cool weather during spring and fall (65°-80°F) or in early morning or late evening for best results. Treat new mounds as they appear. If possible, make applications after a recent rainfall.


VPG

Voluntary Purchasing Groups, Inc.

230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 7401-454
EPA Est. No. 7401-TX-01
Visit Us At: www.hi-yield.com
Product Questions? 855-270-4776

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

32220-0615-WP



2
32221 32220
7