



ferti·lome®

Weed-Out®

Lawn Weed Killer

Contains TRIMEC® Herbicide

- Kills troublesome broadleaf weeds: Dandelion, Chickweed, Clover, Henbit & many others

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See Back Panel For Additional
Precautionary Statements and First Aid

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid 5.45%
 Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 5.88%
 Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) 1.21%

OTHER INGREDIENTS:..... 87.46%

TOTAL: 100.00%

See back panel for substatement of ingredient statement.

NET CONTENTS: ONE GALLON (128 FL. OZS. / 3.785 LITERS)

fertiome® Weed-Out® Lawn Weed Killer

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If In Eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15–20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If On Skin Or Clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15–20 minutes.• Call a poison control center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

In the event of a medical or chemical emergency, contact  North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585.

Manufactured By:



230 FM 87
BONHAM, TX 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-TX-01
Visit Us At: www.fertilo.com
Product Questions? 855-270-4776



7 32221 10519 5

PEEL BACK BOOK HERE

BROADLEAF WEEDS

Annual Yellow Sweetclover	Buttonweed	Fiddleneck	Narrowleaf Vetch	Scarlet Pimpernel	Virginia Creeper
Aster	Canada Thistle	Florida Pusley	Nettle	Scotch Thistle	Virginia Pepperweed
Austrian Fieldcress	Carolina Geranium	Frenchweed	Orange Hawkweed	Sheep Sorrel	Wavyleaf
Bedstraw	Carpetweed	Galinsoga	Oriental Cocklebur	Sheeperspurse	Bullthistle
Beggarticks	Catchweed	Goldenrod	Oxalis	Slender Plantain	Western Clematis
Betony, Florida	Bedstraw	Ground Ivy	Parsley-Piert	Smallflower	Western Salsify
Bindweed, Field	Catnip	Gumweed	Parsnip	Galinsoga	White Mustard
Bird Vetch	Catsear	Hairy Fleabane	Pearlwort	Smooth Dock	Wild Aster
Bitter Wintercress	Chickweed,	Hawkweed	Pennycress	Smooth Pigweed	Wild Buckwheat
Bittercress, Hairy	Common	Healall	Pennywort	Sorrel	Wild Carrot
Bitterweed	Chickweed,	Heartleaf Drymary	Pepperweed	Sowthistle	Wild Four-O'Clock
Black-Eyed Susan	Mouseear	Heathaster	Pigweed	Spanishneedles	Wild Garlic
Black Medic	Chicory	Hedge Bindweed	Pineywoods Bedstraw	Speedwell	Wild Geranium
Black Mustard	Cinquefoil	Hedge Mustard	Plains Coreopsis	Spiny Amaranth	Wild Lettuce
Blackseed Plantain	Clover, Crimson	Henbit	Plantain	Spiny Cocklebur	Wild Marigold
Blessed Thistle	Clover, Hop	Hoary Cress	Poison Ivy	Spiny Sowthistle	Wild Mustard
Bloodflower	Clover, Red	Hoary Plantain	Poison Oak	Spotted Catsear	Wild Onion
Milkweed	Clover, Strawberry	Hoary Vervain	Pokeweed	Spotted Spurge	Wild Parsnip
Blue Lettuce	Clover, Sweet	Horsenettle	Poorjoe	Spurweed	Wild Radish
Blue Vervain	Clover, White	Jimsonweed	Prairie Sunflower	Stinging Nettle	Wild Rape
Bracted Plantain	Cockle	Knawel	Prickly Lettuce	Strawberry, India	Wild Strawberry
Bristly Oxtongue	Cocklebur	Knotweed	Prickly Sida	Mock	Wild Sweet Potato
Broadleaf Dock	Common Mullein	Kochia	Prostrate Knotweed	Tall Nettle	Wild Vetch
Broadleaf Plantain	Creeping Jenny	Lambsquarters	Prostrate Pigweed	Tall Vervain	Woodsorrel
Broomweed	Cudweed	Lespedeza	Prostrate Spurge	Tansy Ragwort	Woolly Croton
Buckhorn	Curly Dock	Mallow	Prostrate Vervain	Tansy Mustard	Woolly Morning-glory
Buckhorn Plantain	Daisy, English	Matchweed	Puncturevine	Tanweed	Wormseed
Bulbous Buttercup	Daisy, Fleabane	Mexicanweed	Purslane, Common	Thistle	Yarrow
Bull Thistle	Daisy, Oxeye	Milk Vetch	Ragweed	Trailing	Yellow Rocket
Bullnettle	Dandelion	Morningglory	Red Sorrel	Crownvetch	Yellowflower
Burclover	Dichondra	Mouseear	Redroot Pigweed	Tumble Mustard	Pepperweed
Burdock	Dogfennel	Hawkweed	Redstem Filaree	Tumble Pigweed	
Burning Nettle	Dollarweed	Mugwort	Rough Cinquefoil	Velvetleaf	
Burweed	False Dandelion	Musk Thistle	Rough Fleabane	Venice Mallow	
Buttercup	Falseflax	Mustard	Russian Pigweed	Virginia	
	False Sunflower	Narrowleaf Plantain	Russian Thistle	Buttonweed	

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.

0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

10519-1114-CL



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

KEEP FROM FREEZING.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drain systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
- Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down an indoor or outdoor drain.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Use only lawn sprayer that delivers coarse spray droplets.
- Do not apply as a fine mist because of potential for injury to desirable plants.

- Do not exceed specified dosages for any area.
- Do not spray Carpetgrass, Dichondra, or lawn where desirable clovers are present. Do not apply this product to "Floratam" St. Augustinegrass in Florida.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application.
- Do not spray when air temperatures exceed 85°F.
- After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

PRODUCT INFORMATION

ferti•lome® Weed Out Lawn Weed Killer controls troublesome common lawn weeds, including Poison Ivy, in both Northern and Southern Grasses.

WHEN TO USE

Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating weeds.

BROADCAST TREATMENTS

Limitations on broadcast applications to residential turfgrass: The maximum application rate is 4 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 fl. oz. of product per 1,000 sq. ft.

BROADCAST TREATMENTS WITH HOSE-END SPRAYERS WITH DILUTION:

Spray preparation for hose-end dilution sprayers:

Add the required amount of **ferti•lome® Weed-Out Lawn Weed Killer** and fill sprayer with water to proper level.

RATES OF ferti•lome® Weed Out Lawn Weed Killer FOR VARIOUS LAWN GRASSES					
Gallons Of Water Spray		Area to Treat (Sq. Ft.)	St. Augustinegrass, Centipedegrass (fl. oz.)	Bermudagrass, Buf- falograss Bentgrass (fl. oz)	Bluegrass Zoysiagrass, Fescue (fl. oz)
Hose End Sprayer	Pressure Sprayer				
1	½	250	½	2/3	2/3 to 1
4	2	1,000	2	2 2/3	2 2/3 to 4
8	4	2,000	4	5 1/3	5 1/3 to 8
20	10	5,000	10	13 1/3	13 1/3 to 20
32	16	8,000	16	21 1/3	21 1/3 to 32

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup
8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)
32 fluid ounces = 1 quart (qt.)

SPOT TREATMENT WITH PRESSURE SPRAYERS

Limitations on spot treatments to residential turfgrass: Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4 fl. oz. per 1,000 sq. ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

ST. AUGUSTINEGRASS, CENTIPEDEGRASS, AND BAHIAGRASS: Mix 2/3 fl. oz. (4 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. Do not apply to St. Augustinegrass, Centipedegrass, or Bahiagrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage. Do not use on "Floratam" St. Augustinegrass in Florida.

BERMUDAGRASS, BUFFALOGRASS, AND BENTGRASS: Mix 1 fl. oz. (2 Tbs) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

KENTUCKY BLUEGRASS, FESCUE, PERENNIAL RYEGRASS, AND ZOYSIAGRASS: Mix 1-1/3 fl. oz. (8 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.