



ferti·lome®

**WEED ZONE
FREE**

Ready To Use

- Rain-fast in as little as 3 hours
- Visible results in just 24 hours
- Proven cool-weather performance
- For home lawns only

TRIMEC® is a registered trademark of PBI/Gordon Corporation

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See attached booklet for complete
Precautionary Statements and
Directions for Use, including
Storage and Disposal and First Aid.

ACTIVE INGREDIENTS:

MCPA, 2-ethylhexyl ester	0.3370%
Mecoprop-p acid	0.0660%
Dicamba acid	0.0180%
Carfentrazone-ethyl.....	0.0023%

OTHER INGREDIENTS:.....99.5767%

TOTAL: 100.0000%

THIS PRODUCT CONTAINS: 2-methyl-4-chlorophenoxyacetic acid
equivalent, 0.2160%

NET CONTENTS ONE QUART (32 FL. OZ. / 946ML)

fertiome® Weed Free Zone Ready-To-Use



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first five minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If On Skin or On Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

In the event of a medical or chemical emergency, contact
North America 1-800-255-3924




ENVIRONMENTAL HAZARDS

This pesticide may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 48 hours.

Manufactured By:
*Volumary
Purchasing
Groups, Inc.*



230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 2217-865-7401
EPA Est. No. 7401-TX-01
Visit Us At: www.fertiome.com
Product Questions?
855-270-4776



7 32221 10528 7

PEEL FROM CORNER OF BOOKLET



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.
- Do not apply this product to Bentgrass, Carpetgrass, Dichondra, legumes and lawns where desirable clovers are present.
- Do not apply to 'Floratam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.
- Do not apply more than 1.9 gallons of product per 1,000 sq.ft. per year (1.5 lb a.i. MCPA per acre per year)
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

PRODUCT FACTS:

ferti•lome® WEED FREE ZONE Ready-To-Use contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

ferti•lome® WEED FREE ZONE Ready-To-Use offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in Turfgrass, e.g., Spurge, Pennywort (Dollarweed), Dandelion, and White Clover.
- Injury to weeds can be noticed within hours of application and plant death can occur within 7 - 14 days.

WHERE TO USE

On Lawns: Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as:

Kentucky Bluegrass	Common Bermudagrass	Hybrid Bermudagrass
Perennial Ryegrass	Mixtures of cool season species	Red or Fine Leaf Fescues
Tall Fescue	St. Augustinegrass (excluding 'Floratam')	Zoysiagrass

WHEN TO APPLY

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant Bermudagrass, Zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

HOW TO USE

ferti•lome® WEED FREE ZONE Ready-To-Use is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

BROADLEAF WEEDS CONTROLLED

ferti•lome® WEED FREE ZONE Ready-To-Use will control or suppress the following broadleaf weeds:

BROADLEAF WEEDS

Annual yellow sweetclover	Brassbuttons	Cockle	Fiddleneck
Aster	Bristly oxtongue	Cocklebur	Filaree, redstem
Aster, white prairie	Broomweed	Cocklebur, Oriental	Filaree, whitestem
Aster, heath	Burclover	Cocklebur, spiny	Fleabane, daisy
Aster, wild	Burdock	Common mullein	Fleabane, hairy
Austrian fieldcress	Burweed	Compassplant,	Fleabane, rough
Bedstraw	Buttercup	Creeping Jenny	Florida pusley
Bedstraw, catchweed	Buttercup, bulbous	(Creeping Charlie)	Frenchweed
Bedstraw, creeping	Buttercup,	Cudweed,	Galinsoga
Bedstraw, pinewoods	creeping	Cudweed, purple	Galinsoga, smallflower
Beggarticks	Carolina geranium	Daisy, English	Goathead
Beggarweed,	Carpetweed	Daisy, oxeye	Goldenrod
creeping	Catnip	Dandelion	Ground ivy
Betony, Florida	Chickweed,	Dayflower	Groundsel
Bindweed	common	Dichondra	Gumweed
Bindweed, field	Chickweed,	Dock	Hawkweed
Bindweed, hedge	mouseear	Dock, broadleaf	Hawkweed, orange
Bitter wintercress	Chicory	Dock, curly	Healall
Bittercress, hairy	Cinquefoil	Dock, smooth	Heartleaf drymary
Bitterweed	Cinquefoil rough	Dogbane	Hemp
Black medic,	Clover	Dogfennel	Henbit
Black-eyed Susan	Clover, crimson	Dollarweed	Hoary cress
Bloodflower	Clover, hop	Elderberry	Innocence
milkwweed	Clover, red	False dandelion	(Blue-eyed Mary)
Blue lettuce	Clover, straw	(*spotted catsear	Jimsonweed
	Clover, sweet	& common catsear)	Knawel
	Clover, white	Falseflax	

BROADLEAF WEEDS (Continued)

Knotweed, prostrate	Pearlwort	Purslane, common	Vervain, prostrate
Kochia	Pennsylvania smartweed	Ragweed	Vervain, tall
Lambsquarters	Pennycress	Red sorrel ("sheep sorrel)	Vetch, bird
Lespedeza	Pennywort	Redweed	Vetch, milk
Lespedeza, common	(*dollarweed)	Scarlet pimpernel	Vetch, narrowleaf
Mallow	Peppergrass	Shepherd's purse	Vetch, wild
Mallow, common	(pepperweed)	Sowthistle	Virginia buttonweed
Matchweed	Pigweed	Sowthistle, spiny	Virginia creeper
Mexicanweed	Pigweed, prostrate	Spanish needles	Virginia pepperweed
Morningglory	Pigweed, redroot	Spiny amaranth	Water pennywort
Morningglory, woolly	Pigweed, Russian	Spurge, prostrate	Wavyleaf bullthistle
Mugwort,	Pigweed, smooth	Spurge, spotted	Western clematis
Mustard	Pigweed, tumble	Strawweed	Western safsify
Mustard, black	Pineappleweed	Strawberry,	Wild buckwheat
Mustard, hedge	Plains coreopsis (tickseed)	India mock	Wild carrot
Mustard, tansy	Plantain	Sunflower, false	Wild four-o'clock
Mustard, tumble	Plantain, blackseed	Sunflower, prairie	Wild garlic
Mustard, white	Plantain, bracted	Tansy ragwort	Wild geranium
Mustard, wild	Plantain, broadleaf	Tanweed	Wild lettuce
Nettle	Plantain, buckhorn	Thistle	Wild marigold
Nettle, bull	Plantain, hoary	Thistle, blessed	Wild onion
Nettle, burning	Plantain, narrowleaf	Thistle, bull	Wild parsnip
Nettle, dead	Plantain, slender	Thistle, Canada	Wild radish
Nettle, horse	Plantain, woolly	Thistle, musk	Wild rape
Nettle, stinging	Poison ivy	Thistle, Russian	Wild strawberry
Nettle, tall	Poison oak	Thistle, Scotch	Wild sweet potato
Old world diamond flower	Pokeweed	Trailing crownvetch	Wild violet
Oxalis	Poorjoe	Velvetleaf	Woolly croton
(*yellow woodsorrel & creeping wood sorrel)	Prickly lettuce	Venice mallow	Wormseed
Parsley-piert	Prickly sida	Veronica (*corn speedwell)	Yarrow
Parsnip	Puncturevine	Vervain, blue	Yellow rocket
	Purslane	Vervain, hoary	Yellowflower pepperweed

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in an area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418
EPA Reg. No. 2217-865-7401
EPA Est. No. 7401-TX-01
Visit Us At: www.fertilome.com
Product Questions? 855-270-4776

10528-1-0819-REG

