




Spurge



Crabgrass



Dandelion

ferti·lome®

**Weed-Out**

With **CRABGRASS KILLER**

Ready To Spray

**Controls Crabgrass & Broadleaf Weeds in Lawns**

SEE INSIDE BOOKLET FOR A COMPLETE LIST OF WEEDS CONTROLLED

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE BACK PANEL AND BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt.....	6.42%
Quinclorac .....	2.13%
Dicamba, dimethylamine salt .....	0.60%
<b>OTHER INGREDIENTS:</b> .....	<b>90.85%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>

THIS PRODUCT CONTAINS: 0.456 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%, 0.182 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%, 0.043 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%, Isomer Specific By AOAC Methods.

**NET CONTENTS ONE QUART (32 FL. OZ. / 946 ML)**

ferti•lome® WeedOut with Cr abgrass Killer Ready-To-Spray



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**FIRST AID**

<b>If In Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

In the event of a medical or chemical emergency, contact **ChemTel**   
North America 1-800-255-3924

**ENVIRONMENTAL HAZARDS:**

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

**Manufactured By:**



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 2217-896-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776 7 3 2 2 2 1 1 1 0 3 1 1



PEEL FROM CORNER OF BOOKLET

If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

**Kills Crabgrass, Dandelion, Clover Plus  
200 Other Listed Weeds**

- One Quart Treats 5000 Sq. Ft.
- Kills Even Tough Weeds-Roots & All
- Systemic Herbicide Activity

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 2217-896-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

11031-0914-CL



### **ENVIRONMENTAL HAZARDS (Continued)**

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### **USER SAFETY RECOMMENDATIONS**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **USE RESTRICTIONS:**

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until sprays have dried.

Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.

Do not spray exposed roots of ornamentals and trees.

#### **SHAKE WELL BEFORE USING**

## PRODUCT FACTS

**What It Does:** Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)

**Where to Use:** Lawns: Kills the weeds not the lawn when used as directed.

**When to Use:** Apply when daytime temperatures are below 90°F.

## HOW IT WORKS

**ferti•lome® Weed-Out With Crabgrass Killer Ready To Spray** contains three proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

## WHERE TO USE

**ferti•lome® Weed-Out With Crabgrass Killer Ready To Spray** can be applied to residential lawns with the following turfgrass species:

### LAWNS TO BE TREATED

#### Cool Season Turfgrass

#### Warm Season Turfgrass

Kentucky bluegrass	Bermudagrass <sup>1</sup> , common and hybrid
Ryegrass, perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues (red, hard and chewing)	

<sup>1</sup>Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F

DO NOT apply **ferti•lome® Weed Out With Crabgrass Killer Ready To Spray** to the following:

Bahiagrass	Ornamental plants (flowers, trees, groundcovers, landscape beds, and shrubs)
Bentgrass, colonial and creeping	
Carpetgrass	Seashore paspalum
Centipedegrass	St. Augustinegrass
Dichondra	Turfgrass species that are not listed on this label.
Gardens including vegetables, fruit trees, vines and berries	

### GRASS AND BROADLEAF WEEDS CONTROLLED:

A list is presented below:

#### GRASS WEEDS:

Crabgrass (large and smooth)

Signalgrass (Broadleaf)

Foxtail (green, yellow and giant)

#### BROADLEAF WEEDS:

Annual yellow, sweetclover	milkweed	Carolina geranium	Creeping Jenny
Aster	Blue lettuce	Carpetweed	(Creeping Charlie)
Austrian fieldcress	Blue vervain	Catchweed bedstraw	Cudweed
Bedstraw	Bracted plantain	Catnip	Curly dock
Beggarticks	Brassbuttons	Catsear	Daisy, English
Betony	Bristly oxtongue	Chickweed, common	Daisy fleabane
Florida Bindweed, field	Broadleaf dock	Chickweed, mouseear	Daisy, oxeye
Bird vetch	Broadleaf plantain	Chicory	Dandelion
Bitter wintercress	Broomweed	Cinquefoil	Dichondra
Bittercress, hairy	Buckhorn	Clover, crimson	Dogbane
Bitterweed	Buckhorn plantain	Clover, hop	Dogfennel
Black-eyed Susan	Bulbous buttercup	Clover, red	Dollarweed
Black medic	Bull thistle	Clover, white	False dandelion
Black mustard	Bullnettle	Cockle	Falseflax
Blackseed plantain	Burclover	Cocklebur	False sunflower
Blessed thistle	Burdock	Common mullein	Fiddleneck
Bloodflower	Burning nettle		Florida pusley
	Burweed		Frenchweed
	Buttercup		Galinsoga
	Buttonweed		
	Canada thistle		

Goathead	Oriental	Rough fleabane	Virginia
Goldenrod	cocklebur	Russian	buttonweed
Ground ivy	Oxalis	pigweed	Virginia creeper
Gumweed	Parsley-piert	Russian thistle	Virginia
Hairy fleabane	parsnip	Scarlet	pepperweed
Hawkweed	Pearlwort	pimpernel	Wavyleaf
Healall	Pennycress	Scotch thistle	bullthistle
Heartleaf	Pennywort	Sheep sorrel	Western
drymary	Peppergrass	Shepherd	clematis
Heathaster	Pepperweed	spurse	Western salsify
Hedge bindweed	Pigweed	Slender plantain	White mustard
Hedge mustard	Pineywoods	Smallflower	Wild aster
Hemp	bedstraw	galinsoga	Wild buckwheat
Henbit	Plains coreopsis	Smooth dock	Wild carrot
Hoary cress	(tickseed)	Smooth	Wild four-o'clock
Hoary plantain	Plantain	pigweed	Wild garlic
Hoary vervain	Poison ivy	Sorrel	Wild geranium
Horsenettle	Poison oak	Sowthistle	Wild lettuce
Jimsonweed	Pokeweed	Spanishneedles	Wild marigold
Knawel	Poorjoe	Speedwell	Wild mustard
Knotweed	Prairie	Spiny amaranth	Wild onion
Kochia	sunflower	Spiny cocklebur	Wild parsnip
Lambsquarters	Prickly lettuce	Spiny sowthistle	Wild radish
Lespedeza	Prickly sida	Spotted catsear	Wild rape
Mallow	Prostrate	Spotted spurge	Wild strawberry
Matchweed	knotweed	Spurweed	Wild sweet
Mexicanweed	Prostrate	Stinging nettle	potato
Milk vetch	pigweed	Strawberry,	Wild vetch
Morning glory	Prostrate spurge	India mock	Wild violet
Mouseear	Prostrate	Tall nettle	Woodsorrel
hawkweed	vervain	Tall vervain	Woolly croton
Mugwort	Puncturevine	Tansy ragwort	Woolly morning
Musk thistle	Purslane,	Tansy mustard	glory
Mustard	common	Tanweed	Woolly plantain
Narrowleaf	Ragweed	Thistle	Wormseed
plantain	Red sorrel	Trailing	Yarrow
Narrowleaf	Redroot	crownvetch	Yellow rocket
vetch	pigweed	Tumble mustard	Yellowflower
Nettle	Redstem filaree	Tumble pigweed	pepperweed
Orange	Rough	Velvetleaf	
hawkweed	cinquefoil	Venice mallow	

## WHEN TO USE

Apply this product when weeds are small and actively growing.

Weed Species	Season	Timing	Maximum number of applications <sup>1</sup>	Application interval, days	Comments
<b>Crabgrass</b> (large and smooth):	Spring	From crabgrass emergence until the crabgrass is 3 to 4 inches in height	2	21	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
<b>Foxtail</b> (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	21	Re-treat at the same rate at 21 days later for dense populations.
<b>Broadleaf signal-grass</b>	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	21	Re-treat at the same rate at 21 days later if needed.



<b>Weed Species</b>	<b>Season</b>	<b>Timing</b>	<b>Maximum number of applications<sup>1</sup></b>	<b>Application interval, days</b>	<b>Comments</b>
<b>Dandelion, plantain, but-tercup, speed-well, knotweed</b> and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
<b>Henbit, chick-weed, white clover, ground ivy, lawn burweed</b> and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

#### **WHEN TO USE (Continued)**

<sup>1</sup>The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application

The maximum number of broadcast applications is limited to 2 per year.

A second application may be made 21 days after the initial application.

The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft.

## HOW MUCH TO USE

This bottle will cover the treated areas described when this product is applied at 6.4 fl. oz. of product per 1,000 sq. ft. Over-application or rates above those listed on this label can cause turf injury.

### Instructions:

- Determine the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Attach garden hose to spray nozzle. Turn the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn to be treated. To begin spraying, turn the spray nozzle lever to the "ON" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet. This 32 fl. oz. container will treat 5,000 sq. ft. of turfgrass.
- **FINISH:** To stop spraying, turn spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn spray nozzle to the "Water" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

## STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.