

Crabgrass Preventer Plus Lawn Food Containing Dimension® 11910

# ferti·lome®



## Crabgrass Preventer Plus Lawn Food

Containing  
**Dimension®**

For Control Of Crabgrass And Control Or Suppression  
Of Many Other Annual Grass And Broadleaf Weeds  
In Established Lawns

### 20-0-3

- Can Be Applied Up To 4 Weeks Later Than Other Pre-emergent Crabgrass Preventers
- Contains Dimension® Herbicide
- Feeds & Prevents Crabgrass In One Application

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**


**ACTIVE INGREDIENTS:**

Dithiopyr: 3,5-pyridinedicarbothioic acid,2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester..... 0.125%

**OTHER INGREDIENTS:**..... 99.875%

**TOTAL:** ..... 100.000%

This product contains 0.0625 pound of the active ingredient dithiopyr per 50 pound bag.



NET WEIGHT 32 LBS. (14.5 KG)

**Crabgrass Preventer  
Plus Lawn Food  
Containing Dimension®**

Crabgrass Preventer Plus Lawn Food Containing Dimension®

Crabgrass Preventer Plus Lawn Food Containing Dimension®

**Covers Up To  
10,000 Sq. Ft.**

# Crabgrass Preventer Plus Lawn Food Containing Dimension®

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, do not exceed 4.6 lb. per 1,000 sq. ft. per year of this product (equivalent to 0.25 lb. of active ingredient (ai) per acre).

In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION: Causes eye irritation • Harmful if inhaled**  
Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

## FIRST AID

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**In the event of a medical or chemical emergency contact ChemTel Inc. North America +01-813-248-0585**

## ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

**Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area. **Container Handling:** Nonrefillable Container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## USE DIRECTIONS FOR TURF

This turf product, containing Dimension® herbicide, can be applied up to 4 weeks later than other Crabgrass preventers and provides up to 4 months of Crabgrass control.

**ferti•lome® Crabgrass Preventer Plus Lawn Food Containing Dimension®** provides pre-emergent control of Crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides post-emergent control of seedling Crabgrass after germination and emergence. Post-emergent control is limited to Crabgrass only, but is effective up to 4 weeks after Crabgrass has germinated. This generally corresponds to the time when Crabgrass seedlings first become visible in the established turf grass. Do not apply this product later than 4 weeks after Crabgrass germination.

Florida Applications: Do not exceed 4 applications per year on St. Augustinegrass in central Florida. Do not apply more than 9.2 lb of this product per 1,000 sq ft per application and no more than 27.6 lb of this product/1000 sq ft per year (equivalent to 0.5 and 1.5 lb of active ingredient per acre, respectively).

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed.

COOL-SEASON GRASSES		WARM-SEASON GRASSES	
Bentgrass, Creeping Bluegrass, Kentucky Fescue, Fine Fescue, Tall	Ryegrass, Perennial	Bahiagrass Bermudagrass Buffalograss Carpetgrass	Centipedegrass St. Augustinegrass Zoysiagrass

## USE PRECAUTIONS FOR TURF

This product should be applied only to lawns and turfgrasses that are well established. For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seeding, sodding, or sprigging. Exception: In newly established lawns of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue, this product may be applied two weeks after the first sign of germination.

- Use of this product on lawns and turfgrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfgrasses injury.
- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro aeration, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.
- **Reseeding, Overseeding, or Sprigging:** Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 9.2 lb./1,000 sq. ft. (0.5 lb. active ingredient (ai) per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

## OTHER USE PRECAUTIONS

- Do not use clipping from treated turf for mulching around vegetables or fruit trees
- Do not allow people or pets to enter treated area until dust has settled

## APPLICATION TIMING

### Spring Application

**Control of Crabgrass:** ferti•lome® Crabgrass Preventer Plus Lawn Food Containing Dimension® provides pre-emergent control of Crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides post-emergent control of seedling Crabgrass after germination and emergence. Post-emergent control is limited to Crabgrass only and is effective up to 4 weeks after germination of Crabgrass. This generally corresponds to the time when Crabgrass seedlings are at the 3-leaf stage of growth and first become visible in the established turf. This product will not control Crabgrass at later growth stages when Crabgrass plants begin to tiller (spread). Properly timed applications will control existing Crabgrass and prevent further emergence of Crabgrass seedlings. Do not apply this product later than 4 weeks after Crabgrass germination.

**Control or Suppression of Other Grass and Broadleaf Weeds:** For control or suppression of other grass and broadleaf weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only Crabgrass will be controlled with applications made up to 4 weeks after weed germination.

### Fall Application

This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of Crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turfgrasses within 12 weeks after application.

### Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

### Weeds Controlled or Suppressed

ferti•lome® Crabgrass Preventer Plus Lawn Food Containing Dimension® will provide control of Crabgrass and control or suppression of other listed grass and broadleaf weeds when applied prior their germination.

GRASSES		BROADLEAF WEEDS	
Barley Barnyardgrass Bluegrass, Annual Brome Crabgrass, Large Crabgrass, Smooth Crabgrass, Southern Crowfootgrass <sup>1</sup> Dallisgrass	Goosegrass Foxtail, Green Foxtail, Yellow Kikuyugrass <sup>1</sup> Oats, Wild Ryegrass (Annual & Perennial) Sandbur Smutgrass	Bittercress <sup>1</sup> Carpeweed <sup>1</sup> Chickweed <sup>1</sup> Geranium, Carolina <sup>1</sup> Henbit Knotweed, Prostrate Lespedeza, Common <sup>1</sup> Marestail Medic, Black Mustard Oxalis, Buttercup	Pineappleweed Pig-weed, Redroot Parsley-piert <sup>1</sup> Purslane, Common Rocket, London Shepherdspurse Speedwell, Corn <sup>1</sup> Spurge, Garden Spurge, Prostrate Spurge, Spotted Woodsorrel, Creeping Woodsorrel, Yellow


<sup>1</sup>Suppression only

11910

## Length of Control

ferti•lome® Crabgrass Preventer Plus Lawn Food Containing Dimension® will provide up to 4 months of Crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Refer to the map to determine the region of the U.S. that includes your location. Use the following table to determine the amount of ferti•lome® Crabgrass Preventer Plus Lawn Food Containing Dimension® to apply in your location based on the desired length of weed control or need for post-emergent Crabgrass control.

### Use "Regions" within the United States:



## Recommended Use Rates

Amount of ferti•lome® Crabgrass Preventer Plus Lawn Food Containing Dimension® to Apply (Pounds of product to apply per 1,000 square feet of turfgrass <sup>1</sup> )		
Region of U.S.	----- Length of Crabgrass Control -----	
	3 - 4 months	Early Post-emergent Control
North	3.2 lb/1000 sq. ft. (0.174 lb ai/acre)	4.0 - 9.2 lb/1000 sq. ft. <sup>††</sup> (0.22 - 0.5 lb ai/acre)
South	4.0 - 4.7 lb/1000 sq. ft. <sup>††</sup> (0.22-0.256 lb ai./acre)	4.0 - 9.2 lb/1000 sq. ft. <sup>††</sup> (0.22 - 0.5 lb ai/acre)

<sup>†</sup> 1,000 sq. ft. equals an area 100 ft. X 10 ft. or square area of about 32 ft. on a side  
<sup>††</sup> Use the higher rate in rate range in areas of high weed pressure or for increased length of control

### Maximum Application Rates:

- Do not apply more than 9.2 lb of this product per 1,000 sq. ft. per application and no more than 27.6 lb of this product/1000 sq. ft. per year (equivalent to 0.5 and 1.5 lb of active ingredient (ai) per acre, respectively).
- In the state of New York, do not exceed 4.6 lb per 1000 sq. ft. per year of this product (equivalent to 0.25 lb of active ingredient (ai) per acre).

## APPLICATION INSTRUCTIONS AND EQUIPMENT

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

Spreader	Spreader Settings Lbs. of product per 1,000 sq. ft.		
	3.2 lbs.	4 lbs.	4.7 lbs.
Scotts®/Republic Accugreen (Drop)	4	4 ¼	4 ½
Scotts®/Republic Speedy Green (Broadcast)	3 ¼	3 ¾	4 ¼
ferti•lome® /Earthway Ev-N-Spred (Broadcast)	14	16	17

20-0-3

## GUARANTEED ANALYSIS

Total Nitrogen (N).....	20%
0.50% Ammoniacal Nitrogen	
19.50% Urea Nitrogen	
Soluble Potash (K <sub>2</sub> O).....	3%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
0.05 Water Soluble Copper (Cu)	
Iron (Fe).....	0.10%
0.05 Water Soluble Iron (Fe)	
Manganese (Mn).....	0.05%
0.05 Water Soluble Manganese (Mn)	
Zinc (Zn).....	0.05%
0.05 Water Soluble Zinc (Zn)	

Derived From: Sulfate Ammonia, Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, and Zinc Sulfate.

F370

## TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for a full refund of purchase price paid. To the extent permitted by law otherwise, use by the Buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

## WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

## INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer.

## LIMITATION OF REMEDIES

To the extent permitted by law the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by Buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law in no case shall Seller be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.

Dimension® is a registered trademark of Dow AgroSciences LLC

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

### Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418  
EPA Reg. No. 62719-483-7401  
EPA Est. No. 7401-TX-01  
Visit Us At: [www.fertilome.com](http://www.fertilome.com)  
Product Questions? 855-270-4776

11910-1114-TP

