



ferti·lome[®]

**WEED ZONE
FREE**

Ready-To-Use

- Rain-fast in as little as 3 hours • Visible results in just 24 hours
- Proven cool-weather performance
- For home lawns only

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See attached booklet for complete Precautionary Statements and Directions for Use, including Storage and Disposal and First Aid.

ACTIVE INGREDIENTS:

MCPA, 2-ethylhexyl ester	0.3370%
Mecoprop-p acid	0.0660%
Dicamba acid	0.0180%
Carfentrazone-ethyl	0.0023%

OTHER INGREDIENTS:	99.5767%
TOTAL:	100.0000%

THIS PRODUCT CONTAINS: 2-methyl-4-chlorophenoxyacetic acid equivalent, 0.2160%

NET CONTENTS: ONE GALLON (128 FL. OZ./ 3.78 LITERS)

fertiome® Weed Free Zone Ready-To-Use

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If On Skin or On Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

In the event of a medical or chemical emergency, contact  North America 1-800-255-3924

ENVIRONMENTAL HAZARDS

This pesticide may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 48 hours.

Manufactured By:



230 FM 87 • Bonham, Texas 75418

EPA Reg. No. 2217-865-7401

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776



PEEL BACK BOOK HERE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.
- Do not apply this product to Bentgrass, Carpetgrass, Dichondra, legumes and lawns where desirable clovers are present.
- Do not apply to 'Floratam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact vegetables, flowers or ornamentals as injury may result.
- Do not apply more than 1.9 gallons of product per 1,000 sq. ft. per year. (1.5 lb a.i. MCPA per acre per year)
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

PRODUCT FACTS:

ferti•lome® WEED FREE ZONE Ready-To-Use contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of weed control.

ferti•lome® WEED FREE ZONE Ready-To-Use offers these advantages:

- Excellent postemergent activity with proven performance for broadleaf weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective weed control for common and troublesome weed species in Turfgrass, e.g., Spurge, Pennywort (Dollarweed), Dandelion, and White Clover.
- Injury to weeds can be noticed within hours of application and plant death can occur within 7 - 14 days.

WHERE TO USE

On Lawns: Use to spot treat broadleaf weeds commonly found in cool season and warm season grasses such as:

Kentucky Bluegrass	Common Bermudagrass	Hybrid Bermudagrass
Perennial Ryegrass	Mixtures of cool season species	Red or Fine Leaf Fescues
Tall Fescue	St. Augustinegrass (excluding 'Floratam')	Zoysiagrass

WHEN TO APPLY

For best results, treat weeds in the spring and the fall when weeds are actively growing. However, this product is effective any time during the growing season.

- Do not apply when air temperatures are above 85°F as lawn injury can occur.
- Hard-to-kill weeds may require a repeat application in 3 weeks.
- Do not water lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded lawns until after the second mowing.
- Delay application to newly sodded, plugged, or sprigged lawns until new grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to dormant Bermudagrass, Zoysiagrass, and St. Augustinegrass may be used to control actively growing broadleaf weeds.

HOW TO USE

ferti•lome® WEED FREE ZONE Ready-To-Use is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure lawns.

BROADLEAF WEEDS CONTROLLED

ferti•lome® WEED FREE ZONE Ready-To-Use will control or suppress the following broadleaf weeds:

BROADLEAF WEEDS

Annual yellow sweetclover	Burclover	Common mullein	Filaree, whitestem	Knotweed, prostrate
Aster	Burdock	Compassplant,	Fleabane, daisy	Kochia
Aster, white prairie	Burweed	Creeping Jenny	Fleabane, hairy	Lambsquarters
Aster, heath	Buttercup	(Creeping Charlie)	Fleabane, rough	Lespedeza
Aster, wild	Buttercup, bulbous	Cudweed,	Florida pusley	Lespedeza, common
Austrian fieldcress	Buttercup, creeping	Cudweed, purple	Frenchweed	Mallow
Bedstraw	Carolin geranium	Daisy, English	Galinsoga	Mallow, common
Bedstraw, catchweed	Carpetweed	Daisy, oxeye	Galinsoga,	Matchweed
Bedstraw, pineywoods	Catnip	Dandelion	smallflower	Mexicanweed
Beggarticks	Chickweed, common	Dayflower	Goathead	Morningglory
Beggarweed, creeping	Chickweed, mouseear	Dichondra	Goldenrod	Morningglory, woolly
Betony, Florida	Chicory	Dock	Ground ivy	Mugwort,
Bindweed	Cinquefoil	Dock, broadleaf	Groundsel	Mustard
Bindweed, field	Cinquefoil rough	Dock, curly	Gumweed	Mustard, black
Bindweed, hedge	Clover	Dock, smooth	Hawkweed	Mustard, hedge
Bitter wintercress	Clover, crimson	Dogbane	Hawkweed, orange	Mustard, tansy
Bittercress, hairy	Clover, hop	Dogfennel	Healall	Mustard, tumble
Bitterweed	Clover, red	Dollarweed	Heartleaf drymary	Mustard, white
Black medic,	Clover, strawberry	Elderberry	Hemp	Mustard, wild
Black-eyed Susan	Clover, sweet	False dandelion	Henbit	Nettle
Bloodflower milkweed	Clover, white	(*spotted catsear	Hoary cress	Nettle, bull
Blue lettuce	Cockle	& common catsear)	Innocence	Nettle, burning
Brassbuttons	Cocklebur	Falseflax	(Blue-eyed Mary)	Nettle, dead
Bristly oxtongue	Cocklebur, Oriental	Fiddleneck	Jimsonweed	Nettle, horse
Broomweed	Cocklebur, spiny	Filaree, redstem	Knawel	Nettle, stinging

BROADLEAF WEEDS (Continued)

Nettle, tall	Plantain	Sowthistle, spiny	Vervain, hoary	Wild strawberry
Old world diamond flower	Plantain, blackseed	Spanish needles	Vervain, prostrate	Wild sweet potato
Oxalis	Plantain, bracted	Spiny amaranth	Vervain, tall	Wild violet
(*yellow woodsorrel & creeping wood sorrel)	Plantain, broadleaf	Spurge, prostrate	Vetch, bird	Woolly croton
Parsley-piert	Plantain, buckhorn	Spurge, spotted	Vetch, milk	Wormseed
Parsnip	Plantain, hoary	Spurweed	Vetch, narrowleaf	Yarrow
Pearlwort	Plantain, narrowleaf	Strawberry, India mock	Vetch, wild	Yellow rocket
Pennsylvania smartweed	Plantain, slender	Sunflower, false	Virginia buttonweed	Yellowflower pepperweed
Pennygrass	Plantain, woolly	Sunflower, prairie	Virginia creeper	
Pennywort (*dollarweed)	Poison ivy	Tansy ragwort	Virginia pepperweed	
Peppergrass (pepperweed)	Poison oak	Tanweed	Water pennywort	
Pigweed	Pokeweed	Thistle	Wavyleaf bullthistle	
Pigweed, prostrate	Poorjoe	Thistle, blessed	Western clematis	
Pigweed, redroot	Prickly lettuce	Thistle, bull	Western safsify	
Pigweed, Russian	Puncturevine	Thistle, Canada	Wild buckwheat	
Pigweed, smooth	Purslane	Thistle, musk	Wild carrot	
Pigweed, tumble	Purslane, common	Thistle, Russian	Wild four-o' clock	
Pineappleweed	Ragweed	Thistle, Scotch	Wild garlic	
Plains coreopsis (tickseed)	Red sorrel	Trailing crownvetch	Wild geranium	
	(*sheep sorrel)	Velvetleaf	Wild lettuce	
	Redweed	Venice mallow	Wild marigold	
	Scarlet pimpernel	Veronica	Wild onion	
	Shepherd's purse	(*corn speedwell)	Wild parsnip	
	Sowthistle	Vervain, blue	Wild radish	
			Wild rape	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in an area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 2217-865-7401

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

10532-1-0819-REG

