




Spurge



Crabgrass



Dandelion

ferti·lome[®]

Weed-Out With **CRABGRASS KILLER**

Ready To Spray

**Controls Crabgrass &
Broadleaf Weeds in Lawns**

SEE INSIDE BOOKLET FOR A
COMPLETE LIST OF
WEEDS CONTROLLED

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

SEE BACK PANEL AND BOOKLET
FOR FIRST AID AND ADDITIONAL
PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	6.42%
Quinclorac	2.13%
Dicamba, dimethylamine salt	0.60%
OTHER INGREDIENTS:	90.85%
TOTAL:	100.00%

THIS PRODUCT CONTAINS: 0.456 lb 2,4-dichlorophen-
oxyacetic acid equivalent per gallon or 5.33%, 0.182
lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or
2.13%, 0.043 lb 3,6-dichloro-o-anisic acid equivalent per
gallon or 0.50%, Isomer Specific By AOAC Methods.

511031F-2

NET CONTENTS ONE QUART (32 FL. OZ. / 946 ML)

ferti•lome® Weed-Out™ with Crabgrass Killer Ready-To-Spray



READ THE ENTIRE LABEL FIRST.

OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

In the event of a medical or chemical emergency, contact **ChemTel** 
North America 1-800-255-3924

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Manufactured By:



BONHAM, TEXAS 75418

EPA Reg. No. 2217-896-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776 7 3 2 2 2 1 1 1 0 3 1 1



PEEL FROM CORNER OF BOOKLET

If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

**Kills Crabgrass, Dandelion, Clover Plus
200 Other Listed Weeds**

- One Quart Treats 5000 Sq. Ft.
- Kills Even Tough Weeds-Roots & All
 - Systemic Herbicide Activity

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 2217-896-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

511031B-1



ENVIRONMENTAL HAZARDS

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

SPRAY DRIFT MANAGEMENT

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

SHAKE WELL BEFORE USING

PRODUCT FACTS

What It Does: Kills major broadleaf and troublesome grass weeds including Dandelion, Chickweed and Clover as well as Crabgrass and Foxtail. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed.

When to Use: Apply when daytime temperatures are below 90°F.

HOW IT WORKS

ferti•lome® Weed-Out™ With Crabgrass Killer Ready To Spray contains three proven weed killers that target lawn weeds and Crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE

ferti•lome® Weed-Out™ With Crabgrass Killer Ready To Spray can be applied to residential lawns with the following turfgrass species:

LAWNS TO BE TREATED

Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass	Bermudagrass ¹ , common and hybrid
Ryegrass, perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	

¹Application to Bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F

DO NOT apply **ferti•lome® Weed-Out™ With Crabgrass Killer Ready To Spray** to the following:

Bahiagrass	Ornamental plants (flowers, trees, groundcovers, landscape beds, and shrubs)
Bentgrass, colonial and creeping	
Carpetgrass	Seashore paspalum
Centipedegrass	St. Augustinegrass
Dichondra	Turfgrass species that are not listed on this label.
Gardens including vegetables, fruit trees, vines and berries	

GRASS AND BROADLEAF WEEDS CONTROLLED:

GRASS WEEDS:

Crabgrass (large and smooth)

Signalgrass (Broadleaf)

Foxtail (green, yellow and giant)

BROADLEAF WEEDS:

Annual Yellow,	Milkweed	Carolina	Creeping
Sweetclover	Blue Lettuce	Geranium	Jenny
Aster	Blue Vervain	Carpetweed	(Creeping
Austrian	Bracted Plantain	Catchweed	Charlie)
Fieldcress	Brassbuttons	Bedstraw	Cudweed
Bedstraw	Bristly Oxtongue	Catnip	Curly dock
Beggarticks	Broadleaf Dock	Catsear	Daisy, English
Betony	Broadleaf	Chickweed,	Daisy Fleabane
Florida	Plantain	Common	Daisy, Oxeye
Bindweed, Field	Broomweed	Chickweed,	Dandelion
Bird vetch	Buckhorn	Mouseear	Dichondra
Bitter	Buckhorn	Chicory	Dogbane
Wintercress	Plantain	Cinquefoil	Dogfennel
Bittercress,	Bulbous	Clover, Crimson	Dollarweed
Hairy	Buttercup	Clover, Hop	False Dandelion
Bitterweed	Bull thistle	Clover, Red	Falseflax
Black-eyed	Bullnettle	Clover,	False Sunflower
Susan	Burclover	Strawberry	Fiddleneck
Black Medic	Burdock	Clover, Sweet	Florida Pusley
Black Mustard	Burning Nettle	Clover, White	Frenchweed
Blackseed	Burweed	Cockle	Galinsoga
Plantain	Buttercup	Cocklebur	
Blessed Thistle	Buttonweed	Common	
Bloodflower	Canada Thistle	Mullein	

Goathead	Hawkweed	Cinquefoil	Velvetleaf
Goldenrod	Oriental	Rough Fleabane	Venice Mallow
Ground ivy	Cocklebur	Russian	Virginia
Gumweed	Oxalis	Pigweed	Buttonweed
Hairy Fleabane	Parsley-Piert	Russian Thistle	Virginia Creeper
Hawkweed	Parsnip	Scarlet	Virginia
Healall	Pearlwort	Pimpernel	Pepperweed
Heartleaf	Pennycress	Scotch Thistle	Wavyleaf
Drymary	Pennywort	Sheep Sorrel	Bullthistle
Heathaster	Peppergrass	Shepherd	Western
Hedge Bind	Pepperweed	Spurse	Clematis
weed	Pigweed	Slender Plantain	Western Salsify
Hedge Mustard	Pinewoods	Smallflower	White Mustard
Hemp	Bedstraw	Galinsoga	Wild Aster
Henbit	Plains Coreopsis	Smooth Dock	Wild Buckwheat
Hoary Cress	(tickseed)	Smooth	Wild Carrot
Hoary Plantain	Plantain	Pigweed	Wild Four-o'clock
Hoary Vervain	Poison Ivy	Sorrel	Wild Garlic
Horsenettle	Poison Oak	Sowthistle	Wild Geranium
Jimsonweed	Pokeweed	Spanishneedles	Wild Lettuce
Knawel	Poorjoe	Speedwell	Wild Marigold
Knotweed	Prairie	Spiny Amaranth	Wild Mustard
Kochia	Sunflower	Spiny Cocklebur	Wild Onion
Lambsquarters	Prickly Lettuce	Spiny Sowthistle	Wild Parsnip
Lespedeza	Prickly Sida	Spotted Catsear	Wild Radish
Mallow	Prostrate	Spotted Spurge	Wild Rape
Matchweed	Knotweed	Spurweed	Wild Strawberry
Mexicanweed	Prostrate	Stinging Nettle	Wild Sweet
Milk Vetch	Pigweed	Strawberry,	Potato
Morning glory	Prostrate spurge	India Mock	Wild Vetch
Mouseear	Prostrate	Tall Nettle	Wild Violet
Hawkweed	Vervain	Tall Vervain	Woodsorrel
Mugwort	Puncturevine	Tansy Ragwort	Woolly Croton
Musk thistle	Purslane,	Tansy Mustard	Woolly Morning
Mustard	Common	Tanweed	Glory
Narrowleaf	Ragweed	Thistle	Woolly Plantain
Plantain	Red Sorrel	Trailing	Wormseed
Narrowleaf	Redroot	Crownvetch	Yarrow
Vetch	Pigweed	Tumble Mustard	Yellow Rocket
Nettle	Redstem Filaree	Tumble Pig-	Yellowflower
Orange	Rough	weed	Pepperweed

WHEN TO USE

Apply this product when weeds are small and actively growing.

Weed Species	Season	Timing	Maximum number of applications ¹	Application interval, days	Comments
Crabgrass (large and smooth):	spring	From Crabgrass emergence until the Crabgrass is 3 to 4 inches in height	2	21	Re-treat at the same rate at 21 days later for dense populations or for continued Crabgrass emergence.
Foxtail (green, giant and yellow):	spring and early summer	During the early growth stages or when Foxtail is 1 to 4 inches in height.	2	21	Re-treat at the same rate at 21 days later for dense populations.
Broadleaf signal-grass	spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	21	Re-treat at the same rate at 21 days later if needed.

Weed Species	Season	Timing	Maximum number of applications ¹	Application interval, days	Comments
Dandelion, Plantain, Buttercup, Speedwell, Knotweed and other listed broadleaf weeds	spring	When broadleaf weeds are small and actively growing.	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, Chickweed, White Clover, Ground Ivy, Lawn Burweed and other listed broadleaf weeds	fall	When broadleaf weeds are young and actively growing	2		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

¹The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application

The maximum number of broadcast applications is limited to 2 per year.

A second application may be made 21 days after the initial application.

The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft.

USE PRECAUTIONS:

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- Certain hard-to-kill weeds may need to be spot treated in 21 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

HOW MUCH TO APPLY

This bottle will cover the treated areas described when this product is applied at 6.4 fl. oz. of product per 1,000 sq. ft. Over-application or rates above those listed on this label can cause turf injury.

Instructions:

- Determine the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT:** Attach garden hose to spray nozzle. Turn the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn to be treated. To begin spraying, turn the spray nozzle lever to the "ON" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.

This 32 fl. oz. container will treat 5,000 sq. ft. of turfgrass.

- **FINISH:** To stop spraying, turn spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn spray nozzle to the "Water" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

Page left blank intentionally.