



ferti·lome[®]

Brush Killer • Stump Killer

This Product Is For Outdoor Residential Use Only

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See Back Panel For Additional
Precautionary Statements and
First Aid. In case of emergency
endangering health or the
environment involving this
product, call 1-800-992-5994

ACTIVE INGREDIENTS:

Triclopyr (3,5,6-trichloro-2-Pyridinyloxyacetic acid),
as the Triethylamine salt 8.8%

OTHER INGREDIENTS:..... 91.2%

TOTAL:..... 100.00%

Contains 6.36% Triclopyr acid - 0.6 lb/gallon

Agricultural Chemical: Do not ship or store
with food, feeds, drugs or clothing.

511484-F1

NET CONTENTS ONE PINT (16 FL.OZS. / 4736 ML)

ferti•lome® Brush Killer Stump Killer

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Wear protective clothing such as a long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material, and shoes and socks when applying this product.

FIRST AID

IN THE EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact 
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 62719-226-7401
EPA Est. No. 7401-TX-01

855-270-4776



7 32221 11484 5


PEEL FROM CORNER OF BOOKLET

Manufactured By:



230 FM 87 • BONHAM, TX 75418
EPA Reg. No. 62719-226-7401
EPA Est. No. 7401-TX-01
855-270-4776

511484-B1



USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all Directions For Use carefully before applying.

GENERAL INFORMATION

ferti•lome® BRUSH KILLER STUMP KILLER is recommended for the control of unwanted woody plants, vines and broadleaf weeds around homes, cabins, fences, walkways, and other non-crop areas.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not use on or around edible food or food crops.
- Avoid contact with eyes or clothing during and after application.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- **Entry Restrictions:** Do not enter or allow others to enter the treated area until sprays have dried.
- Do not spray when the wind is blowing toward desirable plants.
- Do not apply this product directly to, or allow to drift to, grapes, vegetable crops, flowers, ornamentals, or other desirable foliage. Applications should be made when wind is low and there is little chance of spray drift.
- Do not apply with nozzles that produce a fine-droplet spray.
- Measuring utensils and spray equipment should not be reused for purposes other than herbicide application.

APPLICATION METHODS

For best results, apply as a full coverage foliar spray when brush and weeds are fully leafed out and actively growing. Wilting or other symptoms will occur in 1 to 6 weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

FOLIAGE SPRAY

Using a hose-end or tank sprayer, mix 8 to 16 tablespoons (4 to 8 fl. oz.) in enough water to make one gallon mixed spray. For mature or hard-to-control plants (such as Ash, Blackgum, Choke Cherry, Elm, Maple, Oak, Pine, or Winged Elm), or during drought

conditions, use the highest rate of application indicated. Resprouting may occur in the year following treatment when lower rates are used. The addition of a surfactant (spreader/sticker) enhances herbicidal action and improved distribution of the spray on hard-to-wet plant surfaces. Apply at the rate of one gallon mixed spray to approximately 500 sq. ft. of area. Coverage may vary depending on the amount of foliage to be treated. Using coarse spray pattern and low pressure, apply as thorough foliage spray to wet all leaves. See

General Use Precautions and Restrictions.

STUMP TREATMENT

To kill stumps and prevent sprouting, apply the undiluted product with a paint brush or backpack sprayer to completely cover the freshly cut surface. Do not reuse paint brush. Wrap in several layers of newspaper and discard in trash.

VINE CONTROL

When treating vines such as Poison Oak, Poison Ivy, etc. that grow on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat as directed for stump treatment.

WOODY PLANTS AND VINES CONTROLLED INCLUDE:

Alder	Douglas Fir	Raspberry
Arkansas Rose	Dogwood	Salmonberry
Arrowwood	Elderberry	Sassafras
Ash	Elm	Scotch Broom
Aspen	Hazel	Sumac

Beech	Honeysuckle	Poplar
Birch	Hornbeam	Sweetbay Magnolia
Blackberry	Kudzu	Sweetgum
Blackgum	Locust	Sycamore
Box Elder	Madrone	Tanoak
Brazilian Pepper	Maples	Thimbleberry
California Rose	Mesquite	Trumpet Creeper
Cascara	Mimosa	Tulip Poplar
Ceanothus	Mulberry	Virginia Creeper
Cherry	Oaks	Western Hemlock
Chinquapin	Persimmon	Willow
Choke Cherry	Pine	Winged Elm
Cottonwood	Poison Ivy	Wild Grape
Crataegus (Hawthorn)	Poison Oak	Wild Rose

PERENNIAL BROADLEAF WEEDS CONTROLLED INCLUDE:

Bindweed	Field Bindweed	Smartweed
Burdock	Field Horsetail Rush	Spurge
Canada Thistle	Henbit	Tansy Ragwort
Chicory	Lambsquarter	Vetch
Clover	Oxalis	Wild Lettuce
Curly Dock	Plantain	Wild Violet
Dandelion	Ragweed	-

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container and keep out of reach of children. Store in a cool, dry place. Avoid contamination of food and foodstuffs.

Container Handling: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Sellers election, one of the following:

1. Refund of purchase price paid by Buyer or User for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Terms and Conditions of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Page left blank intentionally.