




Spurge



Crabgrass



Dandelion

# ferti·lome<sup>®</sup>

## Weed-Out With **CRABGRASS KILLER**

**Controls Crabgrass &  
Broadleaf Weeds in Lawns**

**SEE INSIDE BOOKLET FOR A  
COMPLETE LIST OF  
WEEDS CONTROLLED**

**KEEP OUT OF REACH  
OF CHILDREN**

**CAUTION**

**SEE BACK PANEL AND BOOKLET  
FOR FIRST AID AND ADDITIONAL  
PRECAUTIONARY STATEMENTS**

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt.....	6.42%
Quinclorac .....	2.13%
Dicamba, dimethylamine salt.....	0.60%
<b>OTHER INGREDIENTS:</b> .....	<b>90.85%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>

THIS PRODUCT CONTAINS: 0.456 lb 2,4-dichlorophen-  
oxyacetic acid equivalent per gallon or 5.33%, 0.182  
lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or  
2.13%, 0.043 lb 3,6-dichloro-o-anisic acid equivalent per  
gallon or 0.50%, Isomer Specific By AOAC Methods.

511030F-2

**NET CONTENTS ONE PINT (16 FL. OZ. / 473 ML)**

ferti•lome® Weed-Out™ with Crabgrass Killer



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**


**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**FIRST AID**

<b>If In Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

In the event of a medical or chemical emergency, contact   
North America 1-800-255-3924

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 2217-896-7401

EPA Est. No. 7401-TX-01

855-270-4776



**PEEL FROM CORNER OF BOOKLET**

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418  
EPA Reg. No. 2217-896-7401  
EPA Est. No. 7401-TX-01  
855-270-4776

511030B-2



7 3 2 2 2 1 1 1 0 3 0 4

### **USER SAFETY RECOMMENDATIONS**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### **USE RESTRICTIONS:**

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

### **SPRAY DRIFT MANAGEMENT:**

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

### **HOW MUCH TO APPLY**

**For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass**

Spot Treatments: Mix 2.5 fl. oz. (5 Tbs) in 1 gallon of water to treat 400 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.

Over-application or rates above those specified on this label can cause turf injury.

### **WHEN TO USE**

Apply this product when weeds are small and actively growing.

#### **WEED SPECIES:**

**Crabgrass** ( large and smooth):

Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height. Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.

**Foxtail** ( green, giant and yellow):

Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at the same rate 21 days later for dense populations.

**Broadleaf signalgrass:**

Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage. Re-treat at the same rate at 21 days later if needed.

**Dandelion, Plantain, Buttercup, Speedwell, Knotweed**  
and other listed broadleaf weeds:

Apply in spring when broadleaf weeds are small and actively growing. Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

**Henbit, Chickweed, White Clover, Ground Ivy, Lawn  
Burweed** and other listed broadleaf weeds:

Apply in fall when broadleaf weeds are young and actively growing. Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

The maximum application rate is 6.4 fl.oz. of product per 1,000 sq. ft. per application. The maximum number of applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 12.8 fl. oz. of product per 1,000 sq.ft.

## HOW TO APPLY

### **Pump Style Sprayers**

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.



### **Dial Style Hose-End Sprayers**

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER.**
2. Close sprayer and set dial to 2.5 oz.
3. Connect sprayer to hose
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

### **Concentrate Plus Water Hose-End Sprayers**

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar.
3. Close sprayer, connect to garden hose, shake well.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

## USE PRECAUTIONS

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

## GRASS AND BROADLEAF WEEDS CONTROLLED:

A list is presented below:

### GRASS WEEDS:

Crabgrass (large and smooth)

Signalgrass (Broadleaf)

Foxtail (green, yellow and giant)

### BROADLEAF WEEDS:

Annual Yellow, Sweetclover	Bloodflower Milkweed	Buttercup Buttonweed	Cockle Cocklebur
Aster	Blue Lettuce	Canada Thistle	Common Mullein
Austrian Fieldcress	Blue Vervain Bracted	Geranium	Creeping Jenny
Bedstraw	Plantain	Carpetweed	(Creeping Charlie)
Beggarticks	Brassbuttons	Catchweed Bedstraw	Cudweed
Betony Florida	Bristly Oxtongue	Catnip	Curly dock
Bindweed, Field	Broadleaf Dock Broadleaf	Catsear	Daisy, English
Bird vetch	Plantain	Chickweed, Common	Daisy Fleabane
Bitter Wintercress	Broomweed	Chickweed, Mouseear	Daisy, Oxeye
Bittercress, Hairy	Buckhorn	Chicory	Dandelion
Bitterweed	Buckhorn Plantain	Cinquefoil	Dichondra
Black-eyed Susan	Bulbous Buttercup	Clover,	Dogbane
Black Medic	Bull thistle	Crimson	Dogfennel
Black Mustard	Bullnettle	Clover, Hop	Dollarweed
Blackseed	Burclover	Clover, Red	False Dandelion
Plantain	Burdock	Clover, Strawberry	Falseflax
Blessed Thistle	Burning Nettle	Clover, Sweet	
	Burweed	Clover, White	

False	Matchweed	Poison Ivy	Scarlet
Sunflower	Mexicanweed	Poison Oak	Pimpernel
Fiddleneck	Milk Vetch	Pokeweed	Scotch Thistle
Florida Pusley	Morning glory	Poorjoe	Sheep Sorrel
Frenchweed	Mouseear	Prairie	Shepherd
Galinsoga	Hawkweed	Sunflower	Spurse
Goathead	Mugwort	Prickly Lettuce	Slender
Goldenrod	Musk thistle	Prickly Sida	Plantain
Ground ivy	Mustard	Prostrate	Smallflower
Gumweed	Narrowleaf	Knotweed	Galinsoga
Hairy Fleabane	Plantain	Prostrate	Smooth Dock
Hawkweed	Narrowleaf	Pigweed	Smooth
Healall	Vetch	Prostrate	Pigweed
Heartleaf	Nettle	spurge	Sorrel
Drymary	Orange	Prostrate	Sowthistle
Heathaster	Hawkweed	Vervain	Spanish
Hedge Bind	Oriental	Puncturevine	needles
weed	Cocklebur	Purslane,	Speedwell
Hedge	Oxalis	Common	Spiny
Mustard	Parsley-Piert	Ragweed	Amaranth
Hemp	Parsnip	Red Sorrel	Spiny
Henbit	Pearlwort	Redroot	Cocklebur
Hoary Cress	Pennycress	Pigweed	Spiny
Hoary Plantain	Pennywort	Redstem	Sowthistle
Hoary Vervain	Peppergrass	Filaree	Spotted
Horsenettle	Pepperweed	Rough	Catsear
Jimsonweed	Pigweed	Cinquefoil	Spotted
Knawel	Pineywoods	Rough	Spurge
Knotweed	Bedstraw	Fleabane	Spurweed
Kochia	Plains	Russian	
Lambsquarters	Coreopsis	Pigweed	
Lespedeza	(tickseed)	Russian	
Mallow	Plantain	Thistle	

Stinging Nettle	Virginia	Wild	Woodsorrel
Strawberry,	Buttonweed	Four-o'clock	Woolly Croton
India Mock	Virginia	Wild Garlic	Woolly
Tall Nettle	Creeper	Wild Geranium	Morning
Tall Vervain	Virginia	Wild Lettuce	Glory
Tansy Ragwort	Pepperweed	Wild Marigold	Woolly
Tansy Mustard	Wavyleaf	Wild Mustard	Plantain
Tanweed	Bulbthistle	Wild Onion	Wormseed
Thistle	Western	Wild Parsnip	Yarrow
Trailing	Clematis	Wild Radish	Yellow Rocket
Crownvetch	Western	Wild Rape	Yellowflower
Tumble	Salsify	Wild	Pepperweed
Mustard	White Mustard	Strawberry	
Tumble	Wild Aster	Wild Sweet	
Pigweed	Wild	Potato	
Velvetleaf	Buckwheat	Wild Vetch	
Venice Mallow	Wild Carrot	Wild Violet	

## STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container.  
**If empty:** Place in trash or offer for recycling if available.  
**If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.