

SULFUR

GROUP

M02

FUNGICIDE

TO OPEN...  
CUT HERE!

RESEALABLE  
ZIPPER  
BAG

34188

# Hi-Yield®



## DUSTING WETTABLE SULFUR



- Controls Black Spot, Powdery Mildew, Leaf Spot, Rust & Brown Canker on Roses
- Controls Listed Diseases and Insects on Shrubs, Flowers, Vegetables & Fruits
- Controls Thrips, Rust Mites, Red Spider Mites and Two Spotted Mites on Citrus
- For Use in Residential Lawns and Gardens

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT:

Sulfur ..... 90.0%

INERT INGREDIENTS: ..... 10.0%

TOTAL: ..... 100.0%

### CAUTION

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS



34188

NET WEIGHT 3.25 LBS. (1.47 KG)

# Hi-Yield®

## STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep out of reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container.** Do not reuse this container. **If Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

# Hi-Yield®



## DUSTING WETTABLE SULFUR

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated areas for at least 24 hours and until this material has washed into the soil. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes and being grown in home lawns and gardens.

#### 1. WHAT IS THIS PACKAGE FOR?

This product provides effective and economical control of listed plant diseases and insects. May be used as a dust or spray.

#### 2. HOW DO YOU USE IT?

When used as a dust, apply to all plant surfaces to thoroughly cover.

When used as a spray, mix 4 tablespoons per each 1 gallon of water used. Spray to cover thoroughly all plant surfaces.

**NOTE: DO NOT RE-ENTER TREATED AREAS FOR AT LEAST 24 HOURS AFTER APPLICATION IS MADE.**

### ROSES, SHRUBS, FLOWERS: (as a Fungicide & Insecticide)

For Black Spot, Brown Canker, Leaf Spot, Powdery Mildew, Thrips and Mites on Roses. Begin spraying when new growth starts in the spring. One gallon covers 35 sq. ft. Repeat applications at 7 to 10 day intervals and re-apply after rainfall. For disease and insect control on shrubs, flowers and ornamentals begin application when diseases appear and repeat at 5-10 day intervals and after rains.

For control of Powdery Mildew, Thrips and Mites on Azalea, Alder, Buttonbush, Catalpa, Cosmos, Crape Myrtle, Dahlia, Daisy, Delphinium, Aster, Calendula, Dogwood, Euonymus, Gladiolus, Golden Fleece, Hibiscus, Holly, Hydrangeas, Lady's Mantel, Ligustrum, Lilac, Petunia, Poplars, Silver Vine, Spires, Sunflower, Sweet Pea, Verbena, Violets, Willows, Carnations, Cedar, Cherry Laurel, Chrysanthemums, Hollyhock, Juniper, Simlax and Spruce.

For control of Leaf Spot, Thrips, and Mites on Calendula, Cedar, Cherry Laurel, Chrysanthemum, Clematis, Columbine, Foxglove, Hollyhock, English Ivy, Juniper, Laurel, Sage, Simlax, Snapdragons, and Spruce.

For control of Rust, Thrips, and Mites on Asters, Carnations, Chrysanthemums, Hollyhock and Snapdragon.

### VEGETABLES: (as Fungicide)

For control of Powdery Mildew on Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower. For control of Rust on Beans and Peas. One gallon covers 100 to 1,000 sq. ft. Apply one gallon per 100 sq. ft. during periods of frequent and heavy rainfall. Begin applications at first sign of disease. Repeat applications at 7-10 day intervals. NOTE: Sulfur may injure certain varieties of beans. For control of Powdery and Downy Mildew and Botrytis Blight on Onions. Begin application before diseases are expected to appear. Repeat up to the day of harvest.

**Restrictions:** Do not use on Cucurbits (Cucumbers, Cantaloupes, Melons, and Squash) or Tung.

### FRUITS: (as Fungicide and Insecticide)

**APPLE, PEAR:** For Powdery Mildew and Scab. Reduce dosage to 1 tablespoon (0.33 oz) per gallon of water if hot weather threatens. One gallon covers 60-360 sq. ft. Apply one gallon per 60 sq. ft. during periods of frequent and heavy rainfall. For disease control on apples apply from pre-pink through cover periods. For Pears apply at pre-bloom, bloom, petal fall and cover periods. Repeat applications every 7 to 14 days as needed up to day of harvest.

**PEACH:** For Brown Rot, Scab. Apply in pink, bloom, shuck, cover and pre-harvest sprays. One gallon covers 75-90 sq. ft. Apply one gallon per 75 sq. ft. during periods of frequent and heavy rainfall. Repeat applications every 10 to 14 days as needed up to day of harvest.

**CHERRY:** For Brown Rot, Leaf Spot. Apply in pink, bloom, petal fall, shuck, cover and post-harvest sprays. One gallon covers 70 to 180 sq. ft. Apply one gallon per 70 sq. ft. during periods of frequent and heavy rainfall. Repeat applications every 7 to 14 days as needed up to day of harvest.

**PLUM, PRUNE:** For Brown Rot, Leaf Spot. Apply at pink, bloom, shuck, cover and pre-harvest sprays up to day of harvest. One gallon covers 75-90 sq. ft. Apply one gallon per 75 sq. ft. during periods of frequent and heavy rainfall. Repeat applications every 7 to 14 days as needed up to day of harvest.

**CITRUS:** For Brown Rot. Apply according to state directions up to day of harvest. For Rust, Mite Thrips, Red Spider Mites, Two-spotted Spider Mites. Begin applications at first signs of damage and repeat as needed. Rust Mites should be controlled at all times of the year in order to prevent russetting of fruit and greasy spots on leaves.

**Restriction:** Do not use within 2 weeks of an oil spray treatment, on citrus within 3 to 6 weeks. Use only according to state directions. Sulfur may burn foliage when temperature is high (90°- 95° F).

#### 3. HERE ARE THE RESULTS YOU MAY EXPECT!

When used as directed, this product will provide effective and economical control of listed diseases and insects and thus promote healthy, lush growth.

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 7401-188-34911

EPA Est. No. 7401-TX-01(A); 42403-TX-001 (B)

Letter in lot number corresponds to letter in

parenthesis at the end of the EPA Est. No.

Product Questions? 855-270-4776

34188-101023-REG1




# Hi-Yield®

## FIRST AID

<b>If Inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a Poison Control Center or doctor for treatment advice.</li></ul>
<b>If On Skin Or Clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a Poison Control Center or doctor for treatment advice.</li></ul>
<b>If In Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a Poison Control Center or doctor for treatment advice.</li></ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Call a Poison Control Center or doctor immediately for treatment advice.</li><li>• Have the person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the Poison Control Center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

**In the event of a medical or chemical emergency, contact  North America 1-800-255-3924 or Worldwide Intl. +01- 813-248-0585.**

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear long pants, long sleeved shirt, shoes, socks, water resistant gloves, and goggles or face shield. Wash thoroughly after handling and before eating or smoking.