



BEFORE AFTER

ferti·lome[®]

WEED ZONE FREE

Ready To Spray

- Consistently **Fast** - Controls Tough Weeds Like Clover, Plantain, Ground Ivy, Spurge & Other Listed Weeds
- Proven Cool Weather Performance

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester.....	10.49%
Mecoprop-p, acid.....	2.66%
Dicamba, acid.....	0.67%
Carfentrazone-ethyl.....	0.54%

OTHER INGREDIENTS:	85.64%
TOTAL:	100.00%

THIS PRODUCT CONTAINS:

0.52 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.96%.
0.20 lb. (+)-R-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 2.66%.
0.05 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.67%.
0.04 lb. Ethyl α ,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro 3-methyl-5-oxo-1 H-1,2,4-triazol-1-yl]-4-fluorobenzene propanoate or 0.54%

Isomer Specific by AOAC Methods.

Contains Petroleum Distillate

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See attached booklet for complete Precautionary Statements and Directions for Use, including Storage and Disposal and First Aid.

510527F-1

NET CONTENTS ONE QUART (32 FL. OZ. / 946 ML)



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves.

Personal Hygiene Statements:


After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

FIRST AID

If Swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Do not induce vomiting unless told to by a Poison Control Center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If On Skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate—vomiting may cause aspiration pneumonia.

In the event of a medical or chemical emergency, contact ChemTel  North America 1-800-255-3924

Manufactured By:



230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 2217-855-7401
EPA Est. No. 7401-TX-01
Product Questions? 855-270-4776



PEEL FROM CORNER OF BOOKLET

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 2217-855-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

510527B-1



ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on carpetgrass, dichondra or lawns containing desirable clovers.
- Do not apply this product to 'Floritam' or Bitterblue' varieties of St. Augustinegrass in Florida.
- Do not apply this product to St. Augustinegrass during spring green-up which is the transition period between dormancy and active growth.
- Do not spray roots of trees and ornamentals.
- Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result.
- Do not exceed specified dosages for any area. Be particularly careful within dripline of trees and other ornamental species.
- Do not apply as a fine mist because of potential injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.

- Do not apply when windy. To prevent product runoff, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.

PRODUCT DESCRIPTION

This Trimec® product is a combination of four proven Weed Killers: 2,4-D, mecoprop-p, dicamba and carfentrazone-ethyl. Postemergent applications of this product control a wide range of listed broadleaf Lawn Weeds in Residential Lawns including Dandelion, Chickweed, Spurge, Pennywort (Dollarweed), Knotweed, Plantain, Clover and Ground Ivy (Creeping Charlie).

This product offers these advantages:

- Rapid and effective Weed control for common and troublesome Weed species as listed in turf grass.
- Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed within hours of the application and plant death can occur within 7 - 14 days.

WHERE TO USE

This product is intended for use on most Ornamental Lawns including:

COOL SEASON TURFGRASS	WARM SEASON TURFGRASS
Kentucky Bluegrass	Common Bermudagrass
Perennial Ryegrass	Hybrid Bermudagrass
Tall Fescue	Bahiagrass
Red or Fine Leaf Fescues	Zoysiagrass
Colonial Bentgrass	Buffalograss
Mixtures of Cool Season Species	St. Augustinegrass
	Centipedegrass

Turfgrass Tolerance:

The turfgrass tolerance to **ferti•lome® Weed Free Zone Ready-To-Spray** may vary and temporary turfgrass yellowing may occur on St. Augustinegrass and certain varieties of Hybrid Bermudagrass.

ferti•lome® Weed Free Zone Ready-To-Spray may injure certain turfgrass species. To reduce risk of Lawn injury, avoid spraying drought stressed Lawns or when temperatures are above 90°F.

WHEN TO APPLY

Spray when Broadleaf Weeds are young and actively growing for best results.

ferti•lome® Weed Free Zone Ready-To-Spray may be applied in the spring,

summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2-6 week intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations or as a follow-up treatment, or any time broadleaf weeds are susceptible.

CULTURAL TIPS FOR IMPROVED CONTROL

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. Rain or watering within 4 hours after treatment may reduce effectiveness.

Mowing:

- Delay mowing 1 to 2 days before and after the application of this product.

Reseeding Interval:

- Treated areas may be reseeded 2 weeks after application.

HOW MUCH TO USE

BROADCAST TREATMENTS FOR COOL SEASON TURFGRASS

Use rates of **ferti•lome® Weed Free Zone Ready-To-Spray** on cool season turfgrass as broadcast treatments are presented in Table 1.

TABLE 1. RATE RECOMMENDATIONS FOR COOL SEASON TURFGRASS.

Species	Amount of Product	Treated Area, sq. ft.
	fl. oz./1,000 sq. ft.	1.0 qt. (32 fl. oz.) Sprayer Jar or Retail Container
Kentucky Bluegrass	1.5 to 2.0	16,000 to 21,300 sq.ft
Perennial Ryegrass		
Tall Fescue		
Red or Fine Leaf Fescues		
Colonial Bentgrass		
Mixtures of Cool Season Species		

Use rates of **ferti•lome® Weed Free Zone Ready-To-Spray** on warm season turfgrass as broadcast treatments are presented in Table 2.

TABLE 2. RATE RECOMMENDATIONS FOR WARM SEASON TURFGRASS

Species	Amount of Product	Treated Area, sq. ft.
	fl. oz./1,000 sq. ft.	1.0 qt. (32 fl. oz.) Sprayer Jar or Retail Container
Common Bermudagrass	1.0 to 2.0	16,000 to 32,000 sq. ft.
Hybrid Bermudagrass	0.75 to 1.5	21,300 to 43,000 sq.ft.
Bahiagrass		
Zoysiagrass		
Buffalograss		
St. Augustinegrass		
Centipedegrass		

The maximum application rate is 3.0 fl.oz. of product/1,000 sq.ft. per application (0.5 lb 2,4-D acid equivalent per acre per application). The maximum number of broadcast applications is limited to 2 per year. The maximum seasonal rate is 6.0 fl.oz. of product/1,000 sq.ft., (1.0 lb 2,4-D acid equivalent per acre) excluding spot treatments. Limited to 2 broadcast applications per year. Minimum of 30 days between applications is required.

HOW TO APPLY

FOUR STEPS TO APPLY

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
2. Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the OFF position.
TO BEGIN spraying: Turn on water at faucet. Point nozzle toward lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Spray evenly over the measured area.
3. SPRAY: Spray evenly over the measured area. Walk at a steady pace. Refer to the view strip [graduated scale] on the side of the container to guide coverage. The 32 fl.oz. container will treat 16,000 to 43,000 sq.ft. of turfgrass.
4. To STOP spraying: Turn the control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

BROADLEAF WEEDS CONTROLLED

ferti•lome® Weed Free Zone Ready-To-Spray will control or suppress the following Broadleaf Weeds and will control or suppress other Broadleaf Weeds that are susceptible to 2,4-D.

BROADLEAF WEEDS		
Aster, White Heath And White Prairie	(*Creeping Oxeye)	Poison Ivy
Bedstraw	Filaree, Whitestem And Redstem	Poison Oak
Beggarweed, Creeping	Florida Betony	Prostrate Knotweed
Bindweed	Florida Pusley	(*Knotweed)
Black Medic	Ground Ivy	Puncturevine
Broadleaf Plantain	(*Creeping Charlie)	Purple Cudweed
Buckhorn Plantain	Groundsel	Purslane
Bull Thistle	Hawkweed	Ragweed
Burclover	Healall	Redweed
Burdock, Common	Henbit	Red Sorrel (*Sheep Sorrel)
Buttercup, Creeping	Innocence	Shepherdspurse
Carpeweed	(Blue-eyed Mary)	Spotted Spurge
Chickweed, Common	Lambsquarters	Spurge
Chicory	Lawn Burweed	Thistle
Cinquefoil	Lespedeza, Common	Veronica
Clover	Mallow, Common	(*Corn Speedwell)
Compassplant	Matchweed	Virginia Buttonweed
Curly Dock	Mouseear Chickweed	White Clover
Dandelion	Old World Diamond Flower	(*Dutch Clover, Honeysuckle Clover, White Trefoil And Purplewort)
Dayflower	Oxalis (*Yellow Wood sorrel And Creeping Woodsorrel)	Wild Carrot
Deadnettle	Parsley-piert	Wild Garlic
Dock	Pennsylvania	Wild Geranium
Dogfennel	Smartweed	Wild Lettuce
English Daisy	Pennywort	Wild Mustard
False Dandelion	(*Dollarweed)	Wild Onion
(*Spotted Catsear And Common Catsear)	Pepperweed	Wild Strawberry
Field Bindweed	Pigweed	Wild Violet
(*Morningglory And Creeping Jenny)	Pineappleweed	Yarrow
Field Oxeye-daisy	Plantain	Yellow Rocket

*Synonyms

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the Seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.