



ferti·lome®

Weed-Out®

Lawn Weed Killer

Contains TRIMEC® Herbicide

- **Kills troublesome Broadleaf Weeds: Dandelion, Chickweed, Clover, Henbit & Many Others As Listed**

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

ACTIVE INGREDIENTS:

| | |
|---|-------|
| Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid | 5.45% |
| Dimethylamine salt of 2,4-dichlorophenoxyacetic acid | 5.88% |
| Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) | 1.21% |

| | |
|---------------------------------|---------|
| OTHER INGREDIENTS: | 87.46% |
| TOTAL: | 100.00% |

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.
 0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.
 0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.
 Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

510519F-1

NET CONTENTS: ONE GALLON (128 FL. OZS. / 3.785 LITERS)


ferti•lome® Weed-Out™ Lawn Weed Killer

PEEL BACK BOOK HERE

FIRST AID

| | |
|----------------------------------|---|
| If In Eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15–20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If Swallowed | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| If On Skin Or On Clothing | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15–20 minutes.• Call a poison control center or doctor for treatment advice. |
| If Inhaled | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice. |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

In the event of a medical or chemical emergency, contact  North America 1-800-255-3924.

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.

Manufactured By:



230 FM 87
BONHAM, TX 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-TX-01
Product Questions? 855-270-4776



7 32221 10519 5

KEEP FROM FREEZING.**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.****PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Use only lawn sprayer that delivers coarse spray droplets. Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not exceed specified dosages for any area.
- Do not spray Carpetgrass, Dichondra, or lawn where desirable clovers are present. Do not apply this product to "Floratum" St. Augustinegrass in Florida.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application.
- Do not spray when air temperatures exceed 85°F.
- After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- Failure to follow above precautions may result in injury.

PRODUCT INFORMATION

ferti•lome® Weed-Out™ Lawn Weed Killer controls troublesome common lawn weeds, including Poison Ivy, in both Northern and Southern Grasses.

WHEN TO USE

Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating weeds.

BROADCAST TREATMENTS

Limitations on broadcast applications to residential turfgrass: The maximum application rate is 4 fl. oz. of product per 1,000 sq. ft. per application (0.72 lb 2,4-D ae, 0.57 lb MCP-P ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 fl. oz. of product per 1,000 sq. ft. (1.14 lb 2,4-D ae, 1.06 lb MCP-P ae and 0.24 lb dicamba ae per acre).

BROADCAST TREATMENTS WITH HOSE-END SPRAYERS WITH DILUTION:

Spray preparation for hose-end dilution sprayers:

Add the required amount of **ferti•lome® Weed-Out™ Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Table 1. RATES OF ferti•lome® Weed-Out™ Lawn Weed Killer FOR VARIOUS LAWN GRASSES

| Gallons Of Water Spray | Area to Treat (Sq. Ft.) | St. Augustinegrass, Centipede, Bahiagrass (fl. oz.) | Bermudagrass, Buffalograss, Bentgrass (fl. oz.) | Bluegrass Zoysiagrass, Fescue (fl. oz.) |
|------------------------|-------------------------|---|---|---|
| 1 | 250 | ½ | 2/3 | 2/3 to 1 |
| 4 | 1,000 | 2 | 2 2/3 | 2 2/3 to 4 |
| 8 | 2,000 | 4 | 5 1/3 | 5 1/3 to 8 |
| 20 | 5,000 | 10 | 13 1/3 | 13 1/3 to 20 |
| 32 | 8,000 | 16 | 21 1/3 | 21 1/3 to 32 |

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
 2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup
 8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)
 32 fluid ounces = 1 quart (qt.)

SPOT TREATMENT WITH PRESSURE SPRAYERS

Limitations on spot treatments to residential turfgrass: Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4 fl. oz. per 1,000 sq. ft. per application (0.57 lb MCP-P ae acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

ST. AUGUSTINEGRASS, CENTIPEDEGRASS, AND BAHIAGRASS: Mix 2/3 fl. oz. (4 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. Do not apply to St. Augustinegrass, Centipede, or Bahiagrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage. Do not use on "Floratum" St. Augustinegrass in Florida.

BERMUDAGRASS, BUFFALOGRASS, AND BENTGRASS: Mix 1 fl. oz. (2 Tbs) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

KENTUCKY BLUEGRASS, FESCUE, PERENNIAL RYEGRASS, AND ZOYSIAGRASS: Mix 1-1/3 fl. oz. (8 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

BROADLEAF WEEDS

| | | | | | |
|---------------------------|--------------------|---------------------|---------------------|-------------------|----------------------|
| Annual Yellow Sweetclover | Buttonweed | Fiddleneck | Narrowleaf Vetch | Scarlet Pimpernel | Virginia Creeper |
| Aster | Canada Thistle | Florida Pusley | Nettle | Scotch Thistle | Virginia |
| Austrian Fieldcress | Carolina Geranium | Frenchweed | Orange Hawkweed | Sheep Sorrel | Pepperweed |
| Bedstraw | Catchweed | Galinoga | Oriental Cocklebur | Sheeperspurse | Wavyleaf |
| Beggarticks | Bedstraw | Goldenrod | Oxalis | Slender Plantain | Bullthistle |
| Betony, Florida | Catnip | Ground Ivy | Parsley-Piert | Smallflower | Western Clematis |
| Bindweed, Field | Catsear | Gumweed | Parsnip | Galinoga | Western Salsify |
| Bird Vetch | Chickweed, | Hairy Fleabane | Pearlwort | Smooth Dock | White Mustard |
| Bitter Wintercress | Common | Hawkweed | Pennycress | Smooth Pigweed | Wild Aster |
| Bittercress, Hairy | Chickweed, | Healall | Pennywort | Sorrel | Wild Buckwheat |
| Bitterweed | Mouseear | Heartleaf Drymary | Pepperweed | Sowthistle | Wild Carrot |
| Black-Eyed Susan | Chicory | Heathaster | Pigweed | Spanishneedles | Wild Four-O'Clock |
| Black Medic | Cinquefoil | Hedge Bindweed | Pineywoods Bedstraw | Speedwell | Wild Garlic |
| Black Mustard | Clover, Crimson | Hedge Mustard | Plains Coreopsis | Spiny Amaranth | Wild Geranium |
| Blackseed Plantain | Clover, Hop | Henbit | Plantain | Spiny Cocklebur | Wild Lettuce |
| Blessed Thistle | Clover, Red | Hoary Cress | Poison Ivy | Spiny Sowthistle | Wild Marigold |
| Bloodflower | Clover, Strawberry | Hoary Plantain | Poison Oak | Spotted Catsear | Wild Mustard |
| Milkweed | Clover, Sweet | Hoary Vervain | Pokeweed | Spotted Spurge | Wild Onion |
| Blue Lettuce | Clover, White | Horsenettle | Poorjoe | Spurweed | Wild Parsnip |
| Blue Vervain | Cockle | Jimsonweed | Prairie Sunflower | Stinging Nettle | Wild Radish |
| Bracted Plantain | Cocklebur | Knawel | Prickly Lettuce | Strawberry, India | Wild Rape |
| Bristly Oxtongue | Common Mullein | Knotweed | Prickly Sida | Mock | Wild Strawberry |
| Broadleaf Dock | Creeping Jenny | Kochia | Prostrate Knotweed | Tall Nettle | Wild Sweet Potato |
| Broadleaf Plantain | Cudweed | Lambsquarters | Prostrate Pigweed | Tall Vervain | Wild Vetch |
| Broomweed | Curly Dock | Lespedeza | Prostrate Spurge | Tansy Ragwort | Woodsorrel |
| Buckhorn | Daisy, English | Mallow | Prostrate Vervain | Tansy Mustard | Woolly Croton |
| Buckhorn Plantain | Daisy, Fleabane | Matchweed | Puncturevine | Tanweed | Woolly Morning-glory |
| Bulbous Buttercup | Daisy, Oxeye | Mexicanweed | Purslane, Common | Thistle | Woolly Plantain |
| Bull Thistle | Dandelion | Milk Vetch | Ragweed | Trailing | Wormseed |
| Bullnettle | Dichondra | Morningglory | Red Sorrel | Crownvetch | Yarrow |
| Burclover | Dogfennel | Mouseear | Redroot Pigweed | Turnble Mustard | Yellow Rocket |
| Burdock | Dollarweed | Hawkweed | Redstem Filaree | Turnble Pigweed | Yellowflower |
| Burning Nettle | False Dandelion | Mugwort | Rough Cinquefoil | Vetivetaf | Pepperweed |
| Burweed | Falseflax | Musk Thistle | Rough Fleabane | Venice Mallow | |
| Buttercup | False Sunflower | Mustard | Russian Pigweed | Virginia | |
| | | Narrowleaf Plantain | Russian Thistle | Buttonweed | |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-01

Product Questions? 855-270-4776

510519B-1

