

ferti•lome® Weed-Out™ Broadleaf Weed Control 10925

ferti•lome®



Weed-Out™ Broadleaf Weed Control

- Controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and Many Other Broadleaf Weeds as Listed
- For Use on Bahia, Bermuda, Centipede, Zoysia, Kentucky Bluegrass, Perennial Rye, Fescue and Bentgrass Lawns

ACTIVE INGREDIENTS:		THIS PRODUCT CONTAINS:	
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	1.65%	27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.	
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid.....	0.40%	6.6 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.33%.	
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	0.18%	3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.	
OTHER INGREDIENTS:	97.77%	Isomer Specific by AOAC Methods.	
Total:	100.00%	TRIMEC® is a registered trademark of PBI/Gordon Corporation.	

KEEP OUT OF REACH OF CHILDREN

CAUTION

(SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

NET WEIGHT 10 LBS. (4.54 KG)

ferti•lome® Weed-Out™ Broadleaf Weed Control

ferti•lome® Weed-Out™ Broadleaf Weed Control

10925

ferti•lome®

Weed-Out™ Broadleaf Weed Control



ferti•lome®

Weed-Out™


Broadleaf Weed Control

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.

Text	
If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If On Skin Or On Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
<p>Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.</p>	

In the event of a medical or chemical emergency, contact ChemTel™  North America 1-800-255-3924

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not apply this product to St. Augustinegrass, Carpetgrass, Dichondra, ornamentals, or lawns containing desirable clovers.
- Do not exceed specified dosages for any area. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product runoff, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

WHERE TO USE

ferti•lome® WEED-OUT™ BROADLEAF WEED CONTROL is a Trimec® complex, a combination of three proven weed killers – 2, 4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many others found in home lawns. Established turf of Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Bermudagrass, Zoysiagrass, and Perennial Ryegrass may be treated.

HOW MUCH TO APPLY

For Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Perennial Ryegrass, Bermudagrass, and Zoysiagrass: Apply 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application. The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq. ft.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

CALIBRATION INSTRUCTIONS

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area. Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product. **NOTE: The spreader settings listed are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.**

SPREADER	SPREADER SETTINGS Lbs. of product per 1,000 sq. ft.	
	2 lbs.	4lbs.
Scotts®/Republic Accugreen (Drop)	2 ¼	4 ¼
Scotts®/Republic Speedy Green (Broadcast)	2 ¾	3 ¾
ferti•lome® /Earthway Ev-N-Spred (Broadcast)	12	14

THE FOLLOWING BROADLEAF WEEDS ARE CONTROLLED BY THIS PRODUCT:

Annual Yellow Sweetclover	Chickweed, Common	Hoary Cross	Prostrate Knotweed	Tumble Mustard
Aster	Chickweed, Mouseear	Hoary Plantain	Prostrate Pigweed	Tumble Pigweed
Austrian Fieldcress	Chicory	Hoary Vervain	Prostrate Spurge	Velvetleaf
Bedstraw	Cinquefoil	Horsenettle	Prostrate Vervain	Venice Mallow
Beggarticks	Clover, Crimson	Jimsonweed	Puncturevine	Virginia
Betony, Florida	Clover, Hop	Knawel	Purslane, Common	Virginia Creeper
Bindweed, Field	Clover, Red	Knotweed	Ragweed	Virginia
Bird Vetch	Clover, Strawberry	Kochia	Red Sorrel	Virginia
Bitter Wintercress	Clover, Sweet	Lambsquarters	Redroot Pigweed	Pepperweed
Bittercress, Hairy	Clover, White	Lespedeza	Redstem Filaree	Wavyleaf Bullthistle
Bitterweed	Cockle	Mallow	Rough Cinquefoil	Western Clematis
Black-eyed Susan	Cocklebur	Matchweed	Rough Fleabane	Western Salsify
Black Medic	Common Mullein	Mexicanweed	Russian Pigweed	White Mustard
Black Mustard	Creeping Jenny	Milk Vetch	Russian Thistle	Wild Aster
Blackseed Plantain	Cudweed	Morningglory	Scarlet Pimpernel	Wild Buckwheat
Blessed Thistle	Curly Dock	Mouseear	Scotch Thistle	Wild Carrot
Bloodflower	Daisy, English	Hawkweed	Sheep Sorrel	Wild Four-O'Clock
Milkweed	Daisy, Fleabane	Mugwort	Sheepspurge	Wild Garlic
Blue Lettuce	Daisy, Oxeye	Musk Thistle	Slender Plantain	Wild Geranium
Blue Vervain	Dandelion	Mustard	Smallflower	Wild Lettuce
Bracted Plantain	Dandelion	Narrowleaf Plantain	Galinsoga	Wild Marigold
Bristly Oxtongue	Dichondra	Narrowleaf Vetch	Smooth Dock	Wild Mustard
Broadleaf Dock	Dogfennel	Nettle	Smooth Pigweed	Wild Onion
Broadleaf Plantain	Dollarweed	Orange Hawkweed	Sorrel	Wild Parsnip
Broomweed	False Dandelion	Oriental Cocklebur	Sowthistle	Wild Radish
Buckhorn	Falseflax	Oxalis	Spanishneedles	Wild Rape
Buckhorn Plantain	False Sunflower	Parsley-piert	Speedwell	Wild Strawberry
Bulbous Buttercup	Fiddleneck	Parsnip	Spiny Amaranth	Wild Sweet Potato
Bull Thistle	Florida Pusley	Pearlwort	Spiny Cocklebur	Wild Vetch
Bullnettle Burclover	Frenchweed	Pennycress	Spiny Sowthistle	Woodsorrel
Burdock	Galinsoga	Pepperweed	Spotted Catsear	Woolly Croton
Burning Nettle	Goldenrod	Pigweed	Spotted Spurge	Woolly Morningglory
Burweed	Ground Ivy	Pineywoods	Spurweed	Woolly Plantain
Buttercup	Gumweed	Bedstraw	Stinging Nettle	Wormseed
Buttonweed	Hairy Fleabane	Plains Coreopsis	Strawberry, India	Yarrow
Canada Thistle	Hawkweed	Plantain	Mock	Yellow Rocket
Carolina Geranium	Healall	Poison Ivy	Tall Nettle	Yellowflower
Carpetweed	Heartleaf Drymary	Poison Oak	Tall Vervain	Pepperweed
Catchweed Bedstraw	Heath Aster	Pokeweed	Tansy Ragwort	
Catnip	Hedge Bindweed	Poorjoe	Tansy Mustard	
Catsear	Hedge Mustard	Prairie Sunflower	Tanweed	
Chickweed, Common	Henbit	Prickly Lettuce	Thistle	
		Prickly Sida	Trailing Crownvetch	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER: FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Weed-Out® and ferti•lome® are registered trademarks of Voluntary Purchasing Groups, Inc.

10925-1-100932-REG1

10925

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No.: 2217-657-7401

EPA Est. No.: 7401-TX-01

Product Questions? 855-270-4776

ferti•lome®

Weed-Out™

Broadleaf Weed Control

