

13x22.5"

(26.25" 2up Cylinder Size)

13.00"

3.00"

22.50"

FOLD

0.5" NO PRINT except edge to edge color

0.5" NO PRINT except edge to edge color

Natural GUARD by ferti·lome[®]

Humic 40520

COVERS 2,000 SQ. FT.

• For Use On Lawns, Vegetable Gardens, Flower Beds, Indoor Plants and Compost Bins

0.5" NO PRINT except edge to edge color

This Product Not Intended as a Fertilizer Substitute

KEEP OUT OF REACH OF CHILDREN

NET WEIGHT 20 LBS. (9.07 KG)



Guaranteed By:
VPG Voluntary Purchasing Groups, Inc.
 230 FM 87 • BONHAM, TEXAS 75418
 Visit Us At: www.naturalguard.com
 Product Questions? 855-270-4776

40520-1016-TP

CONTAINS NON-PLANT FOOD INGREDIENT
 35% Humic Acid, Derived From Leonardite F370

Spreader	Spreader Settings
Scotts®/Republic Acquegreen (Drop)	10 lbs of product per 1,000 sq. ft.
Scotts®/Republic Speedy Green (Broadcast)	10
ferti·lome®/Earthway Ev-N-Spread (Broadcast)	9 1/2
	11

A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction.

NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.

APPLICATION EQUIPMENT AND INSTRUCTIONS

This product can be applied with a drop or broadcast (rotary) spreader. For best results, apply this product evenly and uniformly over the turfgrass (lawn) area avoiding streaking, skips or overlaps. Before each application, determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the label states to apply 10 lbs. per 1,000 sq. ft. and your area to treat is 2,000 sq. ft., then place 20 lbs. of product in the hopper and set the spreader setting at 10 (per table below) if using an Acquegreen drop spreader.

FOR NEW LAWNS: Apply 10 to 15 lbs. per 1,000 square feet prior to soil preparation. Till or work into the soil before seeding. Water in after seeding of lawn. Poor, sandy soil will require application rates of 20 to 30 lbs. per 1,000 square feet. Reapply in early fall.

FOR FLOWER BEDS AND VEGETABLE GARDENS: Apply 1 to 1 1/2 lbs. per 100 square feet before soil preparation for planting. Incorporate into the top 2" to 4" of soil at time of planting. Water in after application. Use as needed. This is determined by cropping system, environment and soil type. For best results, make a second application in late fall before tilling or turning under plant stubble.

FOR INDOOR AND PATIO CONTAINER GROWN PLANTS: One teaspoon for 9" or smaller pots; one tablespoon for 12" or larger pots. Broadcast on the surface or incorporate in top 1" to 2" of soil. Apply every 3 to 4 months. Water in after application.

FOR MIXING COMPOST: Add 1/2 cup (1/4 lb.) per cubic foot. Mix well into compost and apply to gardens, flower beds and potted plants.

SPREADER SETTINGS

DIRECTIONS FOR USE

FOR ESTABLISHED LAWNS: Apply 10 lbs. per 1,000 square feet of lawn area in early spring and again in late fall along with normal lawn fertilization program. Soils low in organic matter will respond better to application rates of 20 to 30 lbs. per 1,000 square feet. Water in after application.

0.5" NO PRINT except edge to edge color

0.5" NO PRINT except edge to edge color

Natural GUARD by ferti·lome[®]

Humic GRANULAR HUMIC ACID

3.00"

3.00"

22.50"