



ferti·lome[®]

Weed-Out[®]

Lawn Weed Killer

- Contains TRIMEC[®] Herbicide
- Kills troublesome Broadleaf Weeds: Dandelion, Chickweed, Clover, Henbit & Many Others As Listed
- Covers Up To 8,000 Sq. Ft.

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See Back Panel For Additional
Precautionary Statements and First Aid

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	5.45%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	5.88%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	1.21%
OTHER INGREDIENTS:	87.46%
TOTAL:	100.00%

See back panel for substatement of ingredient statement.

NET CONTENTS ONE QUART (32 FL. OZ. / 946 ML)

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If On Skin Or On Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

In the event of a medical or chemical emergency, contact  North America 1-800-255-3924

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.



BONHAM, TEXAS 75418
 EPA Reg. No. 2217-597-7401
 EPA Est. No. 7401-TX-01
 Visit Us At: www.fertilome.com
 Product Questions? 855-270-4776



Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

10515-1215-TC



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

KEEP FROM FREEZING.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Use only lawn sprayer that delivers coarse spray droplets. Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not exceed specified dosages for any area.

- Do not spray Carpetgrass, Dichondra, or lawn where desirable clovers are present. Do not apply this product to "Floritam" St. Augustinegrass in Florida.
- Do not use this product on or near desirable plants, including contact or spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply to newly seeded lawn grasses until well established. Seed can be sown 3 to 4 weeks after application.
- Do not spray when air temperatures exceed 85°F.
- After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- Failure to follow above precautions may result in injury.

PRODUCT INFORMATION

ferti•lome® Weed-Out® Lawn Weed Killer controls troublesome common Lawn Weeds, including Poison Ivy, in both Northern and Southern Grasses.

WHEN TO USE

Spray at any time when weeds are growing. For best results, apply in spring when weeds are small and growing rapidly. Apply in fall to remove late germinating weeds.

BROADCAST TREATMENTS

Limitations on broadcast applications to residential turfgrass:

The maximum application rate is 4 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 fl. oz. of product per 1,000 sq. ft.

BROADCAST TREATMENTS WITH HOSE-END SPRAYERS WITH DILUTION: Spray preparation for hose-end dilution sprayers: Add the required amount of **ferti•lome® Weed-Out® Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Table 1 - RATES OF ferti•lome® Weed-Out® Lawn Weed Killer FOR VARIOUS LAWN GRASSES

Gallons of Water Spray - Hose End Sprayer with Dilution	Area to Treat (Sq. Ft.)	St. Augustine-grass, Centipede-grass, Bahiagrass (fl.oz.)	Bermudagrass, Buffalograss, Bentgrass (fl.oz)	Bluegrass Zoysiagrass, Fescue (fl.oz)
1	250	½	⅔	⅔ to 1
4	1,000	2	2 ⅔	2 ⅔ to 4
8	2,000	4	5 ⅓	5 ⅓ to 8
20	5,000	10	13 ⅓	13 ⅓ to 20
32	8,000	16	21 ⅓	21 ⅓ to 32

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)

2 fluid ounces = 4 tablespoons 4 fluid ounces = ½ cup

8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)

32 fluid ounces = 1 quart (qt.)

SPOT TREATMENT WITH PRESSURE SPRAYERS

Limitations on spot treatments to residential turfgrass: Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 4 fl. oz. per 1,000 sq. ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

ST. AUGUSTINEGRASS, CENTIPEDEGRASS, AND BAHIAGRASS:

Mix 2/3 fl. oz. (4 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. Do not apply to St. Augustinegrass, centipede-grass, or bahiagrass emerging from winter dormancy; temperatures below 45°F within 72 hours of application may cause damage. Do not use on "Floratam" St. Augustinegrass in Florida.

BERMUDAGRASS, BUFFALOGRASS, AND BENTGRASS:

Mix 1 fl. oz. (2 Tbs) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

KENTUCKY BLUEGRASS, FESCUE, PERENNIAL RYEGRASS, AND ZOYSIAGRASS: Mix 1-1/3 fl. oz. (8 teaspoons) in 1 gallon of water and uniformly spray 330 sq. ft. See Table 1 for additional spray preparations.

- BROADLEAF WEEDS -		
Annual Yellow	Burclover	Dandelion
Sweetclover	Burdock	Dichondra
Aster	Burning Nettle	Dogfennel
Austrian Fieldcress	Burweed	Dollarweed
Bedstraw	Buttercup	False Dandelion
Beggarticks	Buttonweed	Falseflax
Betony, Florida	Canada Thistle	False Sunflower
Bindweed, Field	Carolina Geranium	Fiddleneck
Bird Vetch	Carpetweed	Florida Pusley
Bitter Wintercress	Catchweed Bedstraw	Frenchweed
Bittercress, Hairy	Catnip	Galinsoga
Bitterweed	Catsear	Goldenrod
Black-Eyed Susan	Chickweed, Common	Ground Ivy
Black Medic	Chickweed, Mouseear	Gumweed
Black Mustard	Chicory	Hairy Fleabane
Blackseed Plantain	Cinquefoil	Hawkweed
Blessed Thistle	Clover, Crimson	Healall
Bloodflower Milkweed	Clover, Hop	Heartleaf Drymary
Blue Lettuce	Clover, Red	Heathaster
Blue Vervain	Clover Strawberry	Hedge Bindweed
Bracted Plantain	Clover, Sweet	Hedge Mustard
Bristly	Clover, White	Henbit
Oxtongue Broadleaf	Cockle	Hoary Cress
Dock	Cocklebur	Hoary Plantain
Broadleaf Plantain	Common Mullein	Hoary Vervain
Broomweed	Creeping Jenny	Horsenettle
Buckhorn	Cudweed	Jimsonweed
Buckhorn Plantain	Curly Dock	Knawel
Bulbous Buttercup	Daisy, English	Knotweed
Bull Thistle	Daisy Fleabane	Kochia
Bullnettle	Daisy, Oxeye	Lambsquarters

- BROADLEAF WEEDS (Continued) -

Lespedeza	Puncturevine	Tumble Mustard
Mallow	Purslane,	Tumble Pigweed
Matchweed	Common Ragweed	Velvetleaf
Mexicanweed	Red Sorrel	Venice Mallow
Milk Vetch	Redroot Pigweed	Virginia Buttonweed
Morningglory	Redstem Filaree	Virginia Creeper
Mouseear Hawkweed	Rough Cinquefoil	Virginia Pepperweed
Mugwort	Rough Fleabane	Wavyleaf Bullthistle
Musk Thistle	Russian Pigweed	Western Clematis
Mustard	Russian Thistle	Western Salsify
Narrowleaf Plantain	Scarlet Pimpernel	White Mustard
Narrowleaf Vetch	Scotch Thistle	Wild Aster
Nettle	Sheep Sorrel	Wild Buckwheat Wild
Orange Hawkweed	Sherherdspurse	Carrot
Oriental Cocklebur	Slender Plantain	Wild Four-O'Clock
Oxalis	Smallflower Galinsoga	Wild Garlic
Parsley-Piert	Smooth Dock	Wild Geranium
Parsnip	Smooth Pigweed	Wild Lettuce
Pearlwort	Sorrel	Wild Marigold
Pennycress	Sowthistle	Wild Mustard
Pennywort	Spanishneedles	Wild Onion
Pepperweed	Speedwell Spiny	Wild Parsnip
Pigweed	Amaranth	Wild Radish
Pineywoods Bedstraw	Spiny Cocklebur	Wild Rape
Plains Coreopsis	Spiny Sowthistle	Wild Strawberry
Plantain	Spotted Catsear	Wild Sweet Potato
Poison Ivy	Spotted Spurge	Wild Vetch
Poison Oak	Spurweed Stinging	Woodsorrel
Pokeweed	Nettle	Woolly Croton
Poorjoe	Strawberry India Mock	Woolly
Prairie Sunflower	Tall Nettle	Morningglory
Prickly Lettuce	Tall Vervain	Woolly Plantain
Prickly Sida	Tansy Ragwort	Wormseed
Prostrate Knotweed	Tansy Mustard	Yarrow
Prostrate Pigweed	Tanweed	Yellow Rocket
Prostrate Spurge	Thistle	Yellowflower
Prostrate Vervain	Trailing Crownvetch	Pepperweed

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.

0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.

0.09 lb. 3,6-dichloro-*o*-anisic acid equivalent per gallon or 1.00%.
Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

This page left blank intentionally.